



ELSEVIER

Contents lists available at [ScienceDirect](http://www.sciencedirect.com)

Linear Algebra and its Applications

journal homepage: www.elsevier.com/locate/laaOn the composition of q -skew derivations in Banach algebras

Pjek-Hwee Lee*, Cheng-Kai Liu

Department of Mathematics, National Taiwan University, and Center for Theoretical Sciences, Taipei Office, Taipei 106, Taiwan

Department of Mathematics, National Changhua University of Education, Changhua 500, Taiwan

ARTICLE INFO

Article history:

Received 8 December 2010

Accepted 29 December 2010

Available online 1 February 2011

Submitted by C.K. Li

AMS classification:

47B48

46H15

Keywords:

 q -Skew σ -derivations

Banach algebra

Quasi-nilpotent

Radical

ABSTRACT

Let \mathcal{A} be a Banach algebra, let σ be an automorphism of \mathcal{A} , and let d, δ be q -skew σ -derivations of \mathcal{A} . We show that if $d\delta(a)$ is quasi-nilpotent for any $a \in \mathcal{A}$, then $d\delta(a)^3$ lies in the radical of \mathcal{A} for all $a \in \mathcal{A}$. This result simultaneously generalizes the previous results obtained by Brešar and Šemrl [5], Chebotar et al. [7] and Abdelali [1].

© 2011 Elsevier Inc. All rights reserved.

1. Introduction

Let \mathcal{A} be a Banach algebra over the complex field \mathbb{C} . For $a, b \in \mathcal{A}$, let $[a, b] = ab - ba$ be the commutator of a and b . The classical Kleinecke–Širokov theorem [16,21] states that if a and b are elements in \mathcal{A} such that $[b, [b, a]] = 0$, then $[b, a]$ is *quasi-nilpotent*, namely, it has 0 spectral radius. Put another way, if d is the inner derivation defined by $d(x) = [b, x]$ for $x \in \mathcal{A}$ such that $d^2(a) = 0$ for some element $a \in \mathcal{A}$, then $d(a)$ is quasi-nilpotent. This result was extended later to continuous derivations by Mathieu and Murphy [18] and to arbitrary derivations by Thomas [23].

On the other hand, Pták [19] proved that if d is an inner derivation of \mathcal{A} such that $d^2(a)$ is quasi-nilpotent for every $a \in \mathcal{A}$, then $d^2(a)^2$ lies in the radical of \mathcal{A} for all $a \in \mathcal{A}$. Then Turovskii and Shul'man [24] extended this result to arbitrary derivations. In [5] Brešar and Šemrl characterized commuting pairs of continuous derivations d, δ of \mathcal{A} such that $d\delta(a)$ is quasi-nilpotent for any $a \in \mathcal{A}$. In particular,

* Corresponding author.

E-mail addresses: phlee@math.ntu.edu.tw (P.-H. Lee), ckliucc@ncue.edu.tw (C.-K. Liu).

it turns out that $d\delta(a)^3$ must lie in the radical of \mathcal{A} for all $a \in \mathcal{A}$. Later, Chebotar et al. [7] generalized this result by removing the commutativity assumption $d\delta = \delta d$ and proved: If d, δ are continuous derivations of \mathcal{A} such that $d\delta(a)$ is quasi-nilpotent for any $a \in \mathcal{A}$, then $d\delta(a)^3$ lies in the radical of \mathcal{A} for all $a \in \mathcal{A}$. A closely related result was recently proved by Boudi [3] who showed that if d, δ are continuous derivations of \mathcal{A} such that $d\delta(a)$ has finite spectrum for any $a \in \mathcal{A}$, then $d\delta(a)^3$ lies in the socle of \mathcal{A} modulo the radical of \mathcal{A} for all $a \in \mathcal{A}$.

Let σ be an automorphism of \mathcal{A} . By a σ -derivation δ of \mathcal{A} we mean a linear map $\delta : \mathcal{A} \rightarrow \mathcal{A}$ such that $\delta(ab) = \delta(a)b + \sigma(a)\delta(b)$ for all $a, b \in \mathcal{A}$. Clearly, the $1_{\mathcal{A}}$ -derivations are just the ordinary derivations where $1_{\mathcal{A}}$ is the identity automorphism of \mathcal{A} . The σ -derivations are also called *skew derivations*. Among the skew derivations one important class is that of q -skew derivations as introduced in [11]. A σ -derivation δ of \mathcal{A} is called q -skew if $\sigma\delta\sigma^{-1} = q\delta$, where $0 \neq q \in \mathbb{C}$. Clearly, the 1-skew σ -derivations are just the σ -derivations that commute with the basic automorphism σ . The q -skew derivations appear in q -Weyl algebras, enveloping algebras of solvable Lie superalgebras, and coordinate rings of quantum matrices. See [2,8,9,12–14,17] for some recent results concerning the q -skew derivations. In this paper, we shall generalize Brešar–Šemrl theorem to arbitrary q -skew derivations without assumptions on continuity or commutativity. A special case of continuous 1-skew derivations was recently proved by Abdelali [1] under some extra commutativity assumptions.

More precisely, our main result in this paper is as follows:

Theorem 1.1. *Let \mathcal{A} be a complex Banach algebra, let σ be an automorphism of \mathcal{A} and let d, δ be q -skew σ -derivations of \mathcal{A} . If $d\delta(a)$ is quasi-nilpotent for any $a \in \mathcal{A}$, then $d\delta(a)^3$ lies in the radical of \mathcal{A} for all $a \in \mathcal{A}$.*

2. Preliminaries

Recall that the radical of an algebra \mathcal{A} is equal to the intersection of the kernels of all irreducible representations of \mathcal{A} . Thus, in order to prove Theorem 1.1, it suffices to show that $\pi d\delta(a)^3 = 0$ for all $a \in \mathcal{A}$ and for any irreducible representation π of \mathcal{A} .

Let π be an irreducible representation of the Banach algebra \mathcal{A} on a vector space X . That is, π is an algebra homomorphism of \mathcal{A} into $L(X)$, the algebra of all linear operators on X , such that $\{0\}$ and X are the only subspaces invariant with respect to $\pi(\mathcal{A})$. Let x_0 be any fixed nonzero element of X and define

$$\|x\| = \inf\{\|a\| \mid a \in \mathcal{A}, \pi(a)x_0 = x\}$$

for each $x \in X$. Then it can be shown that $\|\cdot\|$ is a complete norm on X and $\|\pi(a)x\| \leq \|a\|\|x\|$ for all $a \in \mathcal{A}$ and all $x \in X$. (See the proof of [20, Theorem 2.2.6].) Thus $\pi(\mathcal{A}) \subseteq B(X)$, the Banach algebra of all linear operators on X which are bounded with respect to the norm $\|\cdot\|$. Hence, $\|\pi(a)\| \leq \|a\|$ for all $a \in \mathcal{A}$ and so $\pi : \mathcal{A} \rightarrow B(X)$ is continuous.

From now on, π will always denote a continuous irreducible representation of the Banach algebra \mathcal{A} on a Banach space X with the norm defined as above for some fixed $x_0 \in X$. (Note that any two such norms are mutually equivalent. See the proof of [20, Theorem 2.2.7]. So $B(X)$ is well defined by any one of such norms.) Following [4] we call an automorphism σ of \mathcal{A} a π -inner automorphism if there exists an invertible $P \in L(X)$ such that $\pi\sigma(a) = P\pi(a)P^{-1}$ for all $a \in \mathcal{A}$. An automorphism not π -inner is called π -outer. Similarly a σ -derivation δ of \mathcal{A} is called π -inner if there exists $Q \in L(X)$ such that $\pi\delta(a) = \pi\sigma(a)Q - Q\pi(a)$ for all $a \in \mathcal{A}$. Otherwise, it is called π -outer. For a σ -derivation δ , if both σ and δ are π -inner, such operators P, P^{-1} and Q are automatically bounded as shown below. The following proposition is a generalization of [22, Corollary 3.6] without continuity assumptions.

Proposition 2.1. *Let σ be an automorphism of \mathcal{A} and let δ be a σ -derivation of \mathcal{A} . Suppose that there exist $P, Q \in L(X)$ with P invertible in $L(X)$ such that $\pi\sigma(a) = P\pi(a)P^{-1}$ and $\pi\delta(a) = \pi\sigma(a)Q - Q\pi(a)$ for all $a \in \mathcal{A}$. Then P, P^{-1} and Q are all bounded.*

Proof. Clearly, if P, P^{-1} and Q are all bounded, then both $\pi\sigma$ and $\pi\delta$ are continuous. First we verify the continuity of $\pi\sigma$ and $\pi\delta$.

Let \mathcal{P} be the kernel of π ; then \mathcal{P} is a closed ideal of \mathcal{A} so that \mathcal{A}/\mathcal{P} is also a normed algebra. If \mathcal{P} has infinite codimension, then $\pi\delta$ is continuous by [6, Lemma 4.1]. So assume that \mathcal{P} has finite codimension. Since σ and δ are both π -inner, we have $\sigma(\mathcal{P}) \subseteq \mathcal{P}$ and $\delta(\mathcal{P}) \subseteq \mathcal{P}$. Thus the map $\pi\delta : \mathcal{A}/\mathcal{P} \rightarrow B(X)$ defined by $\pi\delta(\bar{a}) = \pi\delta(a)$, where $\bar{a} = a + \mathcal{P} \in \mathcal{A}/\mathcal{P}$, is well defined. Now \mathcal{A}/\mathcal{P} is finite-dimensional, and it is well known that any linear map from a finite-dimensional normed space into another normed space must be continuous. Thus, $\pi\delta$ is continuous. Let (a_n) be a sequence in \mathcal{A} with $\lim_{n \rightarrow \infty} a_n = 0$. Then $\lim_{n \rightarrow \infty} \bar{a}_n = \bar{0}$, where $\bar{a}_n = a_n + \mathcal{P}$, and so $\lim_{n \rightarrow \infty} \pi\delta(a_n) = \lim_{n \rightarrow \infty} \pi\delta(\bar{a}_n) = 0$. Hence, $\pi\delta$ is continuous.

Note that $\sigma - 1_{\mathcal{A}}$ is a σ -derivation of \mathcal{A} and is also π -inner since $\pi(\sigma - 1_{\mathcal{A}})(a) = \pi\sigma(a)I - I\pi(a)$ for all $a \in \mathcal{A}$, where I is the identity operator on X . Hence, by the preceding paragraph, $\pi(\sigma - 1_{\mathcal{A}})$ is continuous and then so is $\pi\sigma$.

Now we show that both P and P^{-1} are bounded. Fix a nonzero $x'_0 \in X$ and set $x_0 = P^{-1}x'_0$. Define a complete norm on X by $\|x\| = \inf\{\|a\| \mid a \in \mathcal{A}, \pi(a)x_0 = x\}$ for $x \in X$. Then, for $x \in X$ and $a \in \mathcal{A}$ with $\pi(a)x_0 = x$, we have

$$\|Px\| = \|\pi P(a)x_0\| = \|P\pi(a)P^{-1}x'_0\| = \|\pi\sigma(a)x'_0\| \leq \|\pi\sigma\| \|a\| \|x'_0\|.$$

Taking the infimum over all such a , we obtain $\|Px\| \leq \|\pi\sigma\| \|x'_0\| \|x\|$. Thus P is bounded and the closed graph theorem implies that P^{-1} is also bounded.

It remains to show that Q is bounded. Choose $a_0 \in \mathcal{A}$ such that $Qx_0 = \pi(a_0)x_0$ and set $\delta' = \delta - \text{ad}_{\sigma}(a_0)$, where $\text{ad}_{\sigma}(a_0)$ is the σ -derivation of \mathcal{A} defined by $\text{ad}_{\sigma}(a_0)(a) = \sigma(a)a_0 - a_0a$ for $a \in \mathcal{A}$. Then δ' is also a π -inner σ -derivation of \mathcal{A} . More precisely, $\pi\delta'(a) = \pi\sigma(a)Q' - Q'\pi(a)$ for all $a \in \mathcal{A}$, where $Q' = Q - \pi(a_0)$. Let $x \in X$. Then if $a \in \mathcal{A}$ with $\pi(a)x_0 = x$, we have

$$\|Q'x\| = \|Q'\pi(a)x_0\| = \|\pi\sigma(a)Q'x_0 - \pi\delta'(a)x_0\| = \|\pi\delta'(a)x_0\| \leq \|\pi\delta'\| \|a\| \|x_0\|,$$

since $Q'x_0 = 0$. Taking the infimum over all such a , we obtain $\|Q'x\| \leq \|\pi\delta'\| \|x_0\| \|x\|$. This implies that Q' , and hence Q , is bounded. \square

Two automorphisms σ, τ of \mathcal{A} are called π -dependent if $\sigma\tau^{-1}$ is π -inner, that is, there exists an invertible $P \in L(X)$ such that $\pi\sigma(a) = P\pi\tau(a)P^{-1}$ for all $a \in \mathcal{A}$. Otherwise, they are called π -independent. Clearly, an automorphism τ of \mathcal{A} and the identity automorphism $1_{\mathcal{A}}$ are π -independent if and only if τ is π -outer. Also, it is easy to see that π -dependency is an equivalence relation for the automorphisms of \mathcal{A} .

Let σ be an automorphism of \mathcal{A} and let $\delta_1, \dots, \delta_n$ be σ -derivations of \mathcal{A} . We say that $\delta_1, \dots, \delta_n$ are \mathbb{C} -dependent modulo π -inner σ -derivations if there exist $\alpha_1, \dots, \alpha_n \in \mathbb{C}$, not all zero, such that $\sum_{i=1}^n \alpha_i \delta_i$ is π -inner. Otherwise, they are said to be \mathbb{C} -independent modulo π -inner σ -derivations. Clearly, a σ -derivation is \mathbb{C} -independent modulo π -inner σ -derivations if and only if it is π -outer.

These notions are generalized in [10] to σ -derivations of \mathcal{A} into $L(X)$. Let σ be an automorphism of \mathcal{A} . By a σ -derivations δ' of \mathcal{A} into $L(X)$ we mean a linear map $\delta' : \mathcal{A} \rightarrow L(X)$ such that $\delta'(ab) = \pi\sigma(a)\delta'(b) + \delta'(a)\pi(b)$ for all $a, b \in \mathcal{A}$. A σ -derivation δ' of \mathcal{A} into $L(X)$ is called *inner* if there exists $Q \in L(X)$ such that $\delta'(a) = \pi\sigma(a)Q - Q\pi(a)$ for all $a \in \mathcal{A}$. Otherwise, it is called *outer*. We say that the σ -derivations $\delta'_1, \dots, \delta'_n$ of \mathcal{A} into $L(X)$ are \mathbb{C} -dependent modulo inner σ -derivations if there exist $\alpha_1, \dots, \alpha_n \in \mathbb{C}$, not all zero, such that $\sum_{i=1}^n \alpha_i \delta'_i$ is inner. Otherwise, they are said to be \mathbb{C} -independent modulo inner σ -derivations.

We shall make use frequently of the following basic result concerning density properties of π -outer automorphisms and σ -derivations.

Theorem 2.2 [10, Theorem 3.4]. *Let σ be an automorphism of \mathcal{A} , τ_1, \dots, τ_m pairwise π -independent automorphisms of \mathcal{A} , and let $\delta'_1, \dots, \delta'_n$ be σ -derivations of \mathcal{A} into $L(X)$ that are \mathbb{C} -independent modulo inner σ -derivations. Then for any \mathbb{C} -independent $x_1, \dots, x_k \in X$ and arbitrary $y_{ij}, z_{\ell j} \in X, i = 1, \dots, n,$*

$\ell = 1, \dots, m$ and $j = 1, \dots, k$, there exists $a \in \mathcal{A}$ such that $\delta_i'(a)x_j = y_{ij}$ and $\pi\tau_\ell(a)x_j = z_{\ell j}$ for all i, ℓ and j .

We remark that either m or n can be 0 in the preceding theorem. In the case that $m = 1$, the assumption of pairwise π -independence holds vacuously.

In what follows, we shall write $\tilde{\theta} = \pi\theta : \mathcal{A} \rightarrow B(X)$ for any linear map θ on \mathcal{A} . Let σ be an automorphism of \mathcal{A} . If δ is a σ -derivation of \mathcal{A} , then $\tilde{\delta}$ is a σ -derivation of \mathcal{A} into $B(X)$. Clearly, the σ -derivations $\delta_1, \dots, \delta_n$ of \mathcal{A} are \mathbb{C} -dependent modulo π -inner σ -derivations if, and only if, the σ -derivations $\delta_1, \dots, \delta_n$ of \mathcal{A} into $B(X)$ are \mathbb{C} -dependent modulo inner σ -derivations. Let d, δ be σ -derivations of \mathcal{A} . For $a, b \in \mathcal{A}$, we have

$$d\delta(ab) = \sigma^2(a)d\delta(b) + d\sigma(a)\delta(b) + \sigma\delta(a)d(b) + d\delta(a)b,$$

and so

$$\tilde{d}\tilde{\delta}(ab) = \tilde{\sigma}^2(a)\tilde{d}\tilde{\delta}(b) + \tilde{d}\tilde{\sigma}(a)\tilde{\delta}(b) + \tilde{\sigma}\tilde{\delta}(a)\tilde{d}(b) + \tilde{d}\tilde{\delta}(a)\pi(b). \quad (2.1)$$

3. Reduction to d, δ being both π -inner

In this section, we shall show that the conclusion of Theorem 1.1 holds if at least one of d and δ is π -outer. We divide the situation into three cases.

Case 1. d and δ are both π -outer.

First we claim that d, δ are \mathbb{C} -dependent modulo π -inner σ -derivations. Assume on the contrary that they are \mathbb{C} -independent modulo π -inner σ -derivations and choose a nonzero $x \in X$. By Theorem 2.2, there exists $b \in \mathcal{A}$ such that $\tilde{\delta}(b)x = x$, $\tilde{d}(b)x = 0$ and $\pi(b)x = 0$. If x and $\tilde{d}\tilde{\delta}(b)x$ are \mathbb{C} -independent, then, by Theorem 2.2 again, there exists $a' \in \mathcal{A}$ such that $\tilde{d}(a')x = x$ and $\tilde{\sigma}(a')\tilde{d}\tilde{\delta}(b)x = 0$. But if $\tilde{d}\tilde{\delta}(b)x = \beta x$ for some $\beta \in \mathbb{C}$, then it follows from the quasi-nilpotency of $\tilde{d}\tilde{\delta}(b)$ that $\beta = 0$ and so we have also $\tilde{d}(a')x = x$ and $\tilde{\sigma}(a')\tilde{d}\tilde{\delta}(b)x = 0$ for some $a' \in \mathcal{A}$. Let $a = \sigma^{-1}(a')$; then we have $\tilde{d}\tilde{\delta}(ab)x = x$ by (2.1), contradicting the quasi-nilpotency of $\tilde{d}\tilde{\delta}(ab)$. Therefore, there exist a nonzero $\alpha \in \mathbb{C}$ and $Q \in L(X)$ such that

$$\tilde{d}(a) = \alpha\tilde{\delta}(a) + \tilde{\sigma}(a)Q - Q\pi(a)$$

for all $a \in \mathcal{A}$.

Next we show $\tilde{d} = \alpha\tilde{\delta}$. Suppose first that σ is π -outer. If $Q = 0$, we are done. So assume $Qx \neq 0$ for some $x \in X$. If Qx and x are \mathbb{C} -independent, then Theorem 2.2 guarantees the existence of $b \in \mathcal{A}$ such that $\tilde{d}(b)x = 0$, $\tilde{\sigma}(b)Qx \neq 0$ and $\pi(b)x = 0$. But if $Qx = \beta x$ for some nonzero $\beta \in \mathbb{C}$, we can pick $b \in \mathcal{A}$ so that $\tilde{d}(b)x = 0$, $\tilde{\sigma}(b)x \neq 0$ and $\pi(b)x = 0$. Then $\tilde{\sigma}(b)Qx \neq 0$ either. Suppose next that σ is π -inner, that is, there exists an invertible $P \in L(X)$ such that $\tilde{\sigma}(a) = P\pi(a)P^{-1}$ and so $\tilde{d}(a) = \alpha\tilde{\delta}(a) + P\pi(a)P^{-1}Q - Q\pi(a)$ for all $a \in \mathcal{A}$. If $P^{-1}Q \in \mathbb{C}I$, then $\tilde{d} = \alpha\tilde{\delta}$, as desired. So assume $P^{-1}Q \notin \mathbb{C}I$; then $P^{-1}Qx$ and x are \mathbb{C} -independent for some $x \in X$. By Theorem 2.2, there exists $b \in \mathcal{A}$ such that $\tilde{d}(b)x = 0$, $\pi(b)x = 0$ and $\pi(b)P^{-1}Qx \neq 0$. Hence $\tilde{\sigma}(b)Qx = P\pi(b)P^{-1}Qx \neq 0$ either. Thus we have shown that if $\tilde{d} \neq \alpha\tilde{\delta}$, there exist $x \in X$ and $b \in \mathcal{A}$ such that $\tilde{d}(b)x = 0$, $\pi(b)x = 0$ and $\tilde{\sigma}(b)Qx \neq 0$. Then $\alpha\tilde{\delta}(b)x = \tilde{d}(b)x - (\tilde{\sigma}(b)Q - Q\pi(b))x = -\tilde{\sigma}(b)Qx \neq 0$ and so $\tilde{\delta}(b)x \neq 0$. If $\tilde{d}\tilde{\delta}(b)x$ and $\tilde{\delta}(b)x$ are \mathbb{C} -independent, by Theorem 2.2, we can choose $a' \in \mathcal{A}$ such that $\tilde{\sigma}(a')\tilde{d}\tilde{\delta}(b)x = 0$ and $\tilde{d}(a')\tilde{\delta}(b)x = x$. Obviously we can also do so even if $\tilde{d}\tilde{\delta}(b)x$ and $\tilde{\delta}(b)x$ are \mathbb{C} -dependent. Let $a = \sigma^{-1}(a')$; then $\tilde{d}\tilde{\delta}(ab)x = x$ by (2.1), contradicting the quasi-nilpotency of $\tilde{d}\tilde{\delta}(ab)$. Consequently, we have $\tilde{d} = \alpha\tilde{\delta}$.

Using $\tilde{d} = \alpha\tilde{\delta}$ and $\sigma\delta = q\delta\sigma$, we obtain from (2.1)

$$\tilde{d}\tilde{\delta}(ab) = \alpha\tilde{\sigma}^2(a)\tilde{\delta}^2(b) + \alpha(1+q)\tilde{\sigma}(a)\tilde{\delta}(b) + \alpha\tilde{\delta}^2(a)\pi(b) \quad (3.1)$$

for all $a, b \in \mathcal{A}$. Let $x \in X$ and $x \neq 0$. By Theorem 2.2, there exists $b \in \mathcal{A}$ such that $\tilde{\delta}(b)x = x$ and $\pi(b)x = 0$. Also, there exists $a' \in \mathcal{A}$ such that $\tilde{\delta}(a')x = x$ and $\tilde{\sigma}(a')\tilde{\delta}^2(b)x = 0$. Let $a = \sigma^{-1}(a')$;

then $\widetilde{d\delta}(ab)x = \alpha(1+q)x$, implying that $1+q=0$. Then it follows from (3.1) that $\widetilde{d\delta}(ab) = \alpha\widetilde{\sigma^2}(a)\widetilde{\delta}(b) + \alpha\widetilde{\delta^2}(a)\pi(b) = \widetilde{\sigma^2}(a)\widetilde{d\delta}(b) + \widetilde{d\delta}(a)\pi(b)$ for all $a, b \in \mathcal{A}$, and so $\widetilde{d\delta}$ is a σ^2 -derivation of \mathcal{A} into $B(X)$.

Moreover, $\widetilde{d\delta}$ must be inner; otherwise, for any nonzero $x \in X$, there would exist $a \in \mathcal{A}$ such that $\widetilde{d\delta}(a)x = x$, a contradiction. Let $Q \in L(X)$ such that $\widetilde{d\delta}(a) = \widetilde{\sigma^2}(a)Q - Q\pi(a)$ for all $a \in \mathcal{A}$. If $Q = 0$, then $\widetilde{d\delta} = 0$ and we are done. Assume $Qx \neq 0$ for some $x \in X$. Then σ^2 is π -inner; otherwise there would exist $a \in \mathcal{A}$ such that $\widetilde{\sigma^2}(a)Qx = x$ and $\pi(a)x = 0$, and so $\widetilde{d\delta}(a)x = x$, a contradiction. Hence there exists an invertible $P \in L(X)$ such that $\widetilde{\sigma^2}(a) = P\pi(a)P^{-1}$ and so $\widetilde{d\delta}(a) = P\pi(a)P^{-1}Q - Q\pi(a)$ for all $a \in \mathcal{A}$. If $P^{-1}Q \notin \mathbb{C}I$, there would exist $x \in X$ such that $P^{-1}Qx, x$ are \mathbb{C} -independent, and $a \in \mathcal{A}$ such that $\pi(a)x = 0$ and $\pi(a)P^{-1}Qx = P^{-1}x$, and then $\widetilde{d\delta}(a)x = x$, a contradiction. Hence, $P^{-1}Q \in \mathbb{C}I$. Thus $\widetilde{d\delta} = 0$ and we are done.

Case 2. d is π -outer and δ is π -inner.

Let $Q \in L(X)$ such that $\widetilde{d}(a) = \widetilde{\sigma}(a)Q - Q\pi(a)$ for all $a \in \mathcal{A}$. We claim $\widetilde{\delta} = 0$ in this case. Assume on the contrary $\widetilde{\delta} \neq 0$; then $Q \neq 0$.

Suppose first that σ is π -outer. Let $x \in X$ with $Qx \neq 0$ and choose $b \in \mathcal{A}$ such that $\widetilde{\sigma}(b)Qx = x$, $\pi(b)x = 0$ and $\widetilde{d}(b)x = 0$. Suppose next that σ is π -inner, say, there exists an invertible $P \in L(X)$ such that $\widetilde{\sigma}(a) = P\pi(a)P^{-1}$ and so $\widetilde{d}(a) = P\pi(a)P^{-1}Q - Q\pi(a)$ for all $a \in \mathcal{A}$. Note that $P^{-1}Q \notin \mathbb{C}I$ since $\widetilde{\delta} \neq 0$, so we can choose $x \in X$ such that $P^{-1}Qx, x$ are \mathbb{C} -independent. Let $b \in \mathcal{A}$ such that $\pi(b)P^{-1}Qx = P^{-1}x$, $\pi(b)x = 0$ and $\widetilde{d}(b)x = 0$; then $\widetilde{\sigma}(b)Qx = x$ holds too. In either situation, we have $\widetilde{\sigma}(b)Qx = x$, $\pi(b)x = 0$ and $\widetilde{d}(b)x = 0$ for some $b \in \mathcal{A}$ and nonzero $x \in X$. Then $\widetilde{\delta}(b)x = x$. Let $a' \in \mathcal{A}$ such that $\widetilde{d}(a')x = x$ and $\widetilde{\sigma}(a')\widetilde{d\delta}(b)x = 0$, and set $a = \sigma^{-1}(a')$; then $\widetilde{d\delta}(ab)x = x$ by (2.1), a contradiction.

Therefore $\widetilde{\delta} = 0$. Then $\widetilde{\sigma\delta} = q\widetilde{\delta}\sigma = 0$ and so it follows from (2.1) that $\widetilde{d\delta}(ab) = \widetilde{\sigma^2}(a)\widetilde{d\delta}(b) + \widetilde{d\delta}(a)\pi(b)$ for all $a, b \in \mathcal{A}$. Thus $\widetilde{d\delta}$ is a σ^2 -derivation of \mathcal{A} into $L(X)$, and hence $\widetilde{d\delta} = 0$ as we have shown in Case 1.

Case 3. d is π -inner and δ is π -outer.

Let $Q \in L(X)$ such that $\widetilde{d}(a) = \widetilde{\sigma}(a)Q - Q\pi(a)$ for all $a \in \mathcal{A}$. Assuming $\widetilde{d} \neq 0$ and proceeding as in Case 2 with d and δ exchanged, we can choose a nonzero $x \in X$ and $b \in \mathcal{A}$ such that $\widetilde{\sigma}(b)Qx = x$, $\pi(b)x = 0$ and $\widetilde{\delta}(b)x = 0$. Then $\widetilde{d}(b)x = x$. Let $a' \in \mathcal{A}$ such that $\widetilde{\delta}(a')x = x$ and $\widetilde{\sigma}(a')\widetilde{d\delta}(b)x = 0$, and set $a = \sigma^{-1}(a')$; then $\widetilde{\sigma\delta}(a)x = q\widetilde{\delta}\sigma(a)x = q\widetilde{\delta}(a')x = qx$ and hence $\widetilde{d\delta}(ab)x = qx$ by (2.1), a contradiction. Hence $\widetilde{d} = 0$. Then $\widetilde{d\delta} = 0$ and we are done.

4. Reduction to σ being π -inner

Thus we may assume henceforth that both d and δ are π -inner. In this section, we shall show that we may assume further that σ is also π -inner. We assume on the contrary that σ is π -outer, and proceed according as whether σ^2 is π -outer. Write $\widetilde{d}(a) = \widetilde{\sigma}(a)Q - Q\pi(a)$ and $\widetilde{\delta}(a) = \widetilde{\sigma}(a)R - R\pi(a)$ for all $a \in \mathcal{A}$, where $Q, R \in L(X)$. Then $\widetilde{d\delta}(a) = \widetilde{\sigma\delta}(a)Q - Q\pi(\delta(a)) = q\widetilde{\delta}(\sigma(a))Q - Q\widetilde{\delta}(a)$, and so

$$\widetilde{d\delta}(a) = q\widetilde{\sigma^2}(a)RQ - qR\widetilde{\sigma}(a)Q - Q\widetilde{\sigma}(a)R + QR\pi(a) \quad (4.1)$$

for all $a \in \mathcal{A}$.

Case 1. σ^2 is π -outer.

Note that $1_{\mathcal{A}}, \sigma, \sigma^2$ are pairwise π -independent in this case. First we have $RQ = 0$; for if $RQx \neq 0$ for some $x \in X$, then by Theorem 2.2, there would exist $a \in \mathcal{A}$ such that $\widetilde{\sigma^2}(a)RQx = x$ and $\widetilde{\sigma}(a)Qx = \widetilde{\sigma}(a)Rx = \pi(a)x = 0$, and so $\widetilde{d\delta}(a)x = qx$ by (4.1), a contradiction. Similarly $QR = 0$; for if $QRx \neq 0$ for some $x \in X$, then by Theorem 2.2, there would exist $a \in \mathcal{A}$ such that $\pi(a)QRx = x$ and $\widetilde{\sigma}(a)Q^2Rx = 0$, and so $\widetilde{d\delta}(a)QRx = QRx$, a contradiction. Thus (4.1) is reduced to

$$\widetilde{d\delta}(a) = -qR\widetilde{\sigma}(a)Q - Q\widetilde{\sigma}(a)R \quad (4.2)$$

for all $a \in \mathcal{A}$.

Now we claim either $Q^2 = 0$ or $R^2 = 0$. Assume on the contrary $Q^2 \neq 0$ and $R^2 \neq 0$. Then we can choose $x \in X$ such that $Q^2x \neq 0$ and $R^2x \neq 0$; otherwise the vector space X would be the set-theoretic union of two proper subspaces, namely the kernels of Q^2 and R^2 . From $QR = RQ = 0$, it follows that Qx and Rx are \mathbb{C} -independent. If Q^2x and R^2x were also \mathbb{C} -independent, there would exist $a \in \mathcal{A}$ such that $\tilde{\sigma}(a)Q^2x = \frac{1}{q}x$ and $\tilde{\sigma}(a)R^2x = x$, and so $\tilde{d}\delta(a)(Qx + Rx) = -(Qx + Rx)$, a contradiction. Hence $R^2x = \alpha Q^2x$ for some nonzero $\alpha \in \mathbb{C}$. Let $\gamma \in \mathbb{C}$ such that $\gamma^2 = \frac{q}{\alpha}$, and $a \in \mathcal{A}$ such that $\tilde{\sigma}(a)Q^2x = x$; then $\tilde{d}\delta(a)(Qx + \gamma Rx) = -\alpha\gamma(Qx + \gamma Rx)$, a contradiction again. Hence either $Q^2 = 0$ or $R^2 = 0$. Thus $\tilde{d}\delta(a)^3 = 0$ for all $a \in \mathcal{A}$ by (4.2) and so we are done.

Case 2. σ^2 is π -inner.

Let $P \in L(X)$ be invertible such that $\tilde{\sigma}^2(a) = P\pi(a)P^{-1}$ for all $a \in \mathcal{A}$. Then (4.1) can be rewritten as

$$\tilde{d}\delta(a) = qP\pi(a)P^{-1}RQ - qR\tilde{\sigma}(a)Q - Q\tilde{\sigma}(a)R + QR\pi(a)$$

for all $a \in \mathcal{A}$. If $P^{-1}RQx$ and x were \mathbb{C} -independent for some $x \in X$, then by Theorem 2.2, there would exist $a \in \mathcal{A}$ such that $\tilde{\sigma}(a)Qx = \tilde{\sigma}(a)Ux = 0$, $\pi(a)x = 0$ and $\pi(a)P^{-1}RQx = P^{-1}x$, and so $\tilde{d}\delta(a)x = qx$, a contradiction. Hence $P^{-1}RQ \in \mathbb{C}I$ and then

$$\tilde{d}\delta(a) = (qRQ + QR)\pi(a) - qR\tilde{\sigma}(a)Q - Q\tilde{\sigma}(a)R$$

for all $a \in \mathcal{A}$. If $y = (qRQ + QR)x \neq 0$ for some $x \in X$, by Theorem 2.2, there would exist $a \in \mathcal{A}$ such that $\pi(a)y = x$ and $\tilde{\sigma}(a)Qy = \tilde{\sigma}(a)Ry = 0$, and so $\tilde{d}\delta(a)y = y$, a contradiction. Hence $qRQ + QR = 0$ and then we have (4.2) again. Thus if $RQ = 0$, then $QR = 0$ and so we are done as in Case 1. Assume $RQ \neq 0$. Then both R and Q are invertible in $L(X)$ since $P^{-1}RQ \in \mathbb{C}I$. If $Q^{-1}Rx$, x were \mathbb{C} -independent for some $x \in X$, then so were Qx and Rx , and there would exist $a \in \mathcal{A}$ such that $\tilde{\sigma}(a)Qx = R^{-1}x$ and $\tilde{\sigma}(a)Rx = 0$, and so $\tilde{d}\delta(a)x = -qx$ by (4.2), a contradiction. Hence $Q^{-1}R = \alpha I$, or equivalently $R = \alpha Q$ for some nonzero $\alpha \in \mathbb{C}$. Thus $\tilde{d}\delta(a) = -\alpha(q+1)Q\tilde{\sigma}(a)Q$ for all $a \in \mathcal{A}$. Choose a nonzero $x \in X$ and $a \in \mathcal{A}$ such that $\tilde{\sigma}(a)Qx = Q^{-1}x$. Then $\tilde{d}\delta(a)x = -\alpha(q+1)x$, implying that $\alpha(q+1) = 0$. So $\tilde{d}\delta = 0$ and we are done.

5. The case d , δ and σ being all π -inner

Now we may assume that d , δ and σ are all π -inner. Thus there exist $P, Q, R \in B(X)$ with P invertible such that $\tilde{\sigma}(a) = P\pi(a)P^{-1}$, $\tilde{d}(a) = \tilde{\sigma}(a)Q - Q\pi(a)$ and $\tilde{\delta}(a) = \tilde{\sigma}(a)R - R\pi(a)$ for all $a \in \mathcal{A}$. We may choose the maps P, Q and R so that Q and R are “ q -skew”-commuting with P in the sense of the following proposition.

Proposition 5.1. *Let σ be an automorphism of \mathcal{A} and let d, δ be q -skew σ -derivations of \mathcal{A} for some nonzero $q \in \mathbb{C}$. Suppose that σ, d, δ are all π -inner. Then there exist $P, Q, R \in B(X)$ with P invertible in $B(X)$ such that $PQP^{-1} = qQ$, $PRP^{-1} = qR$, $\tilde{\sigma}(a) = P\pi(a)P^{-1}$, $\tilde{d}(a) = \tilde{\sigma}(a)Q - Q\pi(a)$ and $\tilde{\delta}(a) = \tilde{\sigma}(a)R - R\pi(a)$ for all $a \in \mathcal{A}$.*

Proof. By Proposition 2.1, there exist $P, Q \in B(X)$ with P invertible in $B(X)$ such that $\pi\sigma(a) = P\pi(a)P^{-1}$ and $\pi d(a) = \pi\sigma(a)Q - Q\pi(a)$ for all $a \in \mathcal{A}$. Then $\pi\sigma^{-1}(a) = P^{-1}\pi(a)P$ for all $a \in \mathcal{A}$. By assumption, we have $\pi\sigma d\sigma^{-1}(a) = q\pi d(a)$ for all $a \in \mathcal{A}$. On the one hand, $\pi\sigma d\sigma^{-1}(a) = P\pi d\sigma^{-1}(a)P^{-1} = P(\pi(a)Q - Q\pi\sigma^{-1}(a))P^{-1} = P(\pi(a)Q - QP^{-1}\pi(a)P)P^{-1} = P(\pi(a)QP^{-1} - QP^{-1}\pi(a))$. On the other hand, $q\pi d(a) = q(\pi\sigma(a)Q - Q\pi(a)) = q(P\pi(a)P^{-1}Q - Q\pi(a)) = qP(\pi(a)P^{-1}Q - P^{-1}Q\pi(a))$. Comparing these two identities, we obtain $\pi(a)(QP^{-1} - qP^{-1}Q) = (QP^{-1} - qP^{-1}Q)\pi(a)$ for all $a \in \mathcal{A}$. Since $\pi(\mathcal{A})$ acts densely on X , it follows that $QP^{-1} - qP^{-1}Q = \lambda I \in \mathbb{C}I$.

If $\lambda = 0$, then $PQP^{-1} = qQ$ as desired. Hence we may assume that $\lambda \neq 0$. We claim that $q \neq 1$ in this case. Assume on the contrary that $q = 1$; then $[Q, P^{-1}] = \lambda I$ and so $[[Q, P^{-1}], P^{-1}] = 0$. Since $Q, P^{-1} \in B(X)$ which is a Banach algebra, it follows from Kleinecke-Shirokov Theorem [16,21] that $[Q, P^{-1}] = \lambda I$ is quasi-nilpotent, a contradiction. Hence $q \neq 1$.

From $PQP^{-1} - qQ = \lambda P$, it follows that $P_1QP_1^{-1} - qQ = P_1$, where $P_1 = \lambda P$. Let $Q_1 = Q + \frac{P_1}{q-1}$; then

$$P_1Q_1P_1^{-1} = P_1QP_1^{-1} + \frac{P_1}{q-1} = qQ + P_1 + \frac{P_1}{q-1} = q\left(Q + \frac{P_1}{q-1}\right) = qQ_1.$$

Moreover, it is easy to see that $\tilde{\sigma}(a) = P_1\pi(a)P_1^{-1}$ and $\tilde{d}(a) = \tilde{\sigma}(a)Q_1 - Q_1\pi(a)$ for all $a \in \mathcal{A}$. Replacing the maps P and Q by P_1 and Q_1 , respectively, we may assume $PQP^{-1} = qQ$ such that $\tilde{\sigma}(a) = P\pi(a)P^{-1}$ and $\tilde{d}(a) = \tilde{\sigma}(a)Q - Q\pi(a)$ for all $a \in \mathcal{A}$. Let $R \in B(X)$ such that $\tilde{\delta}(a) = \tilde{\sigma}(a)R - R\pi(a)$ for all $a \in \mathcal{A}$; then $RP^{-1} - qP^{-1}R = \mu I \in \mathbb{C}I$ as we have seen above. If $\mu = 0$, we are done. Otherwise, $q \neq 1$. Replacing the maps P and R by μP and $R + \frac{\mu P}{q-1}$, respectively, we are done too. \square

Let $P, Q, R \in B(X)$ be as described in Proposition 5.1. Set $A = P^{-1}Q$ and $B = P^{-1}R$; then $PA = qAP$ and $PB = qBP$. Thus, $d(a) = P\pi(a)A - PA\pi(a) = P[\pi(a), A]$, $\tilde{\delta}(a) = P[\pi(a), B]$ and so $d\tilde{\delta}(a) = P[\tilde{\delta}(a), A] = P[P[\pi(a), B], A]$ for all $a \in \mathcal{A}$. Then Theorem 1.1 follows immediately from the following purely algebraic result.

Theorem 5.2. *Let X be a complex vector space and let \mathcal{B} be a dense algebra of linear operators on X . Let A, B, P be linear operators on X with P invertible such that $PA = qAP, PB = qBP$ for some nonzero $q \in \mathbb{C}$. If $H(T) = P[P[T, B], A]$ is quasi-nilpotent for every $T \in \mathcal{B}$, then $H(T)^3 = 0$ for all $T \in \mathcal{B}$.*

Since $PA = qAP, PB = qBP$, an expansion of $H(T)$ yields

$$\begin{aligned} H(T) &= P^2TBA - P^2BTA - PAPTB + PAPBT \\ &= (P^2T)BA - q^2B(P^2T)A - qA(P^2T)B + q^3AB(P^2T). \end{aligned}$$

Thus the hypothesis of Theorem 5.2 can be reformulated as that

$$H_1(T) = TBA - q^2BTA - qATB + q^3ABT \quad (5.1)$$

is quasi-nilpotent for every $T \in \mathcal{B}_1$, where $\mathcal{B}_1 = P^2\mathcal{B}$ is a dense subset of $L(X)$. And, the conclusion $H(T)^3 = 0$ for all $T \in \mathcal{B}$ is just $H_1(T)^3 = 0$ for all $T \in \mathcal{B}_1$. We begin with an interesting consequence of the quasi-nilpotency of $H_1(T)$.

Lemma 5.3. *$BAX \in \text{span}\{Ax, Bx, x\}$ for every $x \in X$. Moreover, for every $x \in X$, there exist $\mu, \nu \in \mathbb{C}$ such that $BAX = \mu Ax + \nu Bx - \mu\nu x$.*

Proof. Assume on the contrary $BAX \notin \text{span}\{Ax, Bx, x\}$ for some $x \in X$. By the density of \mathcal{B}_1 on X , there exists $T \in \mathcal{B}_1$ such that $TBAx = x$ and $TAx = TBx = Tx = 0$. Then $H_1(T)x = x$ by (5.1), contradicting the quasi-nilpotency of $H_1(T)$. Hence, $BAX \in \text{span}\{Ax, Bx, x\}$ for every $x \in X$.

Thus, for each $x \in X$, we may write $BAX = \mu Ax + \nu Bx + \lambda x$ for some $\mu, \nu, \lambda \in \mathbb{C}$. Then it follows from (5.1) that

$$H_1(T)x = (\mu I - q^2B)TAx + (\nu I - qA)TBx + (\lambda I + q^3AB)Tx$$

for all $T \in \mathcal{B}_1$. We proceed to show that μ, ν, λ can be chosen so that $\lambda = -\mu\nu$. Suppose first that Ax, Bx and x are \mathbb{C} -independent. Let $T \in \mathcal{B}_1$ such that $TAx = \nu x, TBx = q^2Bx$ and $Tx = x$; then $H_1(T)x = (\lambda + \mu\nu)x$ and so $\lambda = -\mu\nu$, as desired.

Suppose next that Ax, Bx and x are \mathbb{C} -dependent. If Ax and x are \mathbb{C} -dependent, say $Ax = \alpha x$ for some $\alpha \in \mathbb{C}$, then $BAX = \alpha Bx$ and we are done by taking $\mu = 0$ and $\nu = \alpha$.

So we assume that Ax and x are \mathbb{C} -independent. Suppose first that Bx and x are \mathbb{C} -dependent, say, $Bx = \beta x$ for some $\beta \in \mathbb{C}$. Then by assumption we may write $BAX = \mu Ax + \lambda x$ for some $\mu, \lambda \in \mathbb{C}$

and $ABx = \beta Ax$. So by (5.1)

$$H_1(T)x = (\mu I - q^2 B)TAx + (-\beta qA + \lambda I + q^3 AB)Tx$$

for all $T \in \mathcal{B}_1$. We claim $\lambda = 0$, whence $BAx = \mu Ax$ and so $v = 0$ works. Assume on the contrary $\lambda \neq 0$. Let $T \in \mathcal{B}_1$ such that $TAx = x$ and $Tx = 0$; then $H_1(T)x = (\mu - \beta q^2)x$, implying $\mu = \beta q^2$. Thus $BAx = \beta q^2 Ax + \lambda x$ and

$$H_1(T)x = q^2(\beta I - B)TAx + (-\beta qA + \lambda I + q^3 AB)Tx$$

for all $T \in \mathcal{B}_1$. Now, let $T \in \mathcal{B}_1$ such that $TAx = Ax$ and $Tx = qx$; then $H_1(T)x = \lambda q(1 - q)x$, implying $q = 1$. Thus $BAx = \beta Ax + \lambda x$ and

$$H_1(T)x = (\beta I - B)TAx + (-\beta A + \lambda I + AB)Tx$$

for all $T \in \mathcal{B}_1$. Finally, let $T \in \mathcal{B}_1$ such that $TAx = 0$ and $Tx = x$; then $H_1(T)x = \lambda x$, a contradiction. Hence, $\lambda = 0$ and we are done.

Suppose next that Bx and x are \mathbb{C} -independent. Write $Bx = \alpha Ax + \beta x$ for some $\alpha, \beta \in \mathbb{C}$ with $\alpha \neq 0$. Thus $BAx = \gamma Ax + \theta x$ for some $\gamma, \theta \in \mathbb{C}$. Let $v \in \mathbb{C}$ be a solution of the quadratic equation $\alpha v^2 + (\beta - \gamma)v - \theta = 0$ and set $\mu = \gamma - \alpha v$. Then

$$\begin{aligned} BAx &= (\mu + \alpha v)Ax + (\alpha v^2 + (\beta - \gamma)v)x = \mu Ax + v(\alpha Ax + \beta x) - (\gamma - \alpha v)v x \\ &= \mu Ax + vBx - \mu vx \end{aligned}$$

as required. \square

Lemma 5.4. *Either $H(T)^3 = 0$ for all $T \in \mathcal{B}$ or there exists $y \in X$ such that Ay, By and y are \mathbb{C} -independent.*

Proof. Suppose that Ax, Bx and x are \mathbb{C} -dependent for every $x \in X$. We are going to show that $H(T)^3 = 0$ for all $T \in \mathcal{B}$. According to [5, Theorem 2.4], we have three cases to discuss.

Case 1. A, B and I are \mathbb{C} -dependent.

If $A \in \mathbb{C}I$ or $B \in \mathbb{C}I$, then $H(T) = P[P(T, B), A] = 0$ for all $T \in \mathcal{B}$ and we are done. So we assume that $B = \alpha A + \beta I$ for some $\alpha, \beta \in \mathbb{C}$ with $\alpha \neq 0$. Then $A^2x \in \text{span}\{BAx, Ax\} \subseteq \text{span}\{Ax, Bx, x\} \subseteq \text{span}\{Ax, x\}$ by Lemma 5.3 and so A^2x, Ax and x are \mathbb{C} -dependent for every $x \in X$. By Kaplansky's Theorem [15, Lemma 14], A is algebraic of degree at most 2. So $(A - \gamma_1 I)(A - \gamma_2 I) = 0$ for some $\gamma_1, \gamma_2 \in \mathbb{C}$. Replacing A by $A - \lambda_1 I$, we may assume $A(A - \gamma I) = 0$ for some $\gamma \in \mathbb{C}$. Then

$$\begin{aligned} H(T) &= P[P(T, B), A] = P[P(T, \alpha A), A] \\ &= \alpha(P^2TA^2 - P^2ATA - PAPTA + PAPAT) \\ &= \alpha(\gamma P^2 - P^2A - PAP)TA + \alpha PAPAT = MTA + NT \end{aligned}$$

for all $T \in \mathcal{B}$, where $M = \alpha(\gamma P^2 - P^2A - PAP)$ and $N = \alpha PAP A$. If ANx and Nx were \mathbb{C} -independent for some $x \in X$, there would exist $T \in \mathcal{B}$ such that $TANx = 0$ and $TNx = x$, and so $H(T)Nx = Nx$, a contradiction. Hence $ANx \in \mathbb{C}Nx$ for every $x \in X$. By [7, Lemma 2.7] $AN = \lambda_1 N$ for some $\lambda_1 \in \mathbb{C}$. Similarly, $AM = \lambda_2 M$ and $A(M + N) = \lambda_3(M + N)$ for some $\lambda_2, \lambda_3 \in \mathbb{C}$. Assume that M and N are \mathbb{C} -independent. Then $\lambda_1 = \lambda_2 = \lambda_3$, so we may write $AN = \lambda N$ and $AM = \lambda M$ for some $\lambda \in \mathbb{C}$. Let $x \in X$ such that $(N + \lambda M)x \neq 0$ and let $T \in \mathcal{B}$ such that $T(N + \lambda M)x = x$. Then $H(T)(N + \lambda M)x = (N + \lambda M)x$, a contradiction. Hence $N = \theta M$ for some $\theta \in \mathbb{C}$; then $H(T) = MT(A + \theta I)$ for all $T \in \mathcal{B}$. If $(A + \theta I)Mx \neq 0$ for some $x \in X$, there would exist $T \in \mathcal{B}$ such that $T(A + \theta I)Mx = x$, and so $H(T)Mx = Mx$, a contradiction. Consequently, $(A + \theta I)M = 0$ and then $H(T)^2 = 0$ for all $T \in \mathcal{B}$, as desired.

Case 2. $\dim X \leq 3$.

In this case, $L(X) \cong M_3(\mathbb{C})$, the 3×3 matrix algebra over \mathbb{C} . So $H(T)$ is nilpotent and then $H(T)^3 = 0$ for all $T \in \mathcal{B}$.

Case 3. There is an idempotent $E \in L(X)$ of rank one such that $(I - E)A = \alpha(I - E)$, where $\alpha \in \mathbb{C}$. Then $A - \alpha I = E(A - \alpha I)$ has rank at most one. From

$$H(T) = P[P[T, B], A] = P[P[T, B], A - \alpha I] = P^2[T, B](A - \alpha I) - P(A - \alpha I)P[T, B],$$

it follows that $H(T)$ has rank at most two. As $H(T)$ is quasi-nilpotent, it follows that $H(T)^3 = 0$ for all $T \in \mathcal{B}$. \square

In view of Lemma 5.4, we may assume in what follows that there exists $y \in X$ such that Ay, By and y are \mathbb{C} -independent. Henceforth, we shall fix the notation y for this specified vector. By Lemma 5.3, there exist $\mu, v \in \mathbb{C}$ such that $BAy = \mu Ay + vBy - \mu vy$. We shall also keep the notations μ, v fixed. Set $A' = A - vI$ and $B' = B - \mu I$; then $B'A'y = 0$. Moreover, $A'y, B'y$ and y are also \mathbb{C} -independent and $\text{span}\{A'y, B'y, y\} = \text{span}\{Ay, By, y\}$. Note that it follows from (5.1) that

$$\begin{aligned} H_1(T) &= TB'A' - (q^2B' + \mu(q+1)(q-1)I)TA' - (qA' + v(q-1)I)TB' \\ &\quad + (q^3A'B' + \mu q(q+1)(q-1)A' + vq^2(q-1)B' + \mu v(q+1)(q-1)^2I)T \end{aligned} \quad (5.2)$$

for all $T \in \mathcal{B}_1$.

Lemma 5.5. Suppose that there exists $x \in X$ such that $B'x = 0$ but $B'A'x \neq 0$. Then (1) $v = 0$, (2) $B'A'z = \mu(q+1)(q-1)A'z$ for all $z \in X$ with $B'z = 0$, (3) $A'^2x = 0$.

Proof. By Lemma 5.3, $BAx = \mu_x Ax + v_x Bx - \mu_x v_x x$, or equivalently, $(B - \mu_x I)(A - v_x I)x = 0$ for some $\mu_x, v_x \in \mathbb{C}$. Then $(B' - (\mu_x - \mu)I)(A' - (v_x - v)I)x = 0$ and the expansion of this yields $B'A'x = \alpha(A'x + \beta x)$, where $\alpha = \mu_x - \mu$ and $\beta = -(v_x - v)$. Hence, $\alpha \neq 0$.

Since $B'x = 0$ but $B'A'x \neq 0$, $A'x$ and x are \mathbb{C} -independent. Let $T \in \mathcal{B}_1$ such that $TA'x = x$ and $Tx = 0$; then $H_1(T)x = (\alpha - \mu(q+1)(q-1))x$ by (5.2) and so $\alpha = \mu(q+1)(q-1)$. Thus (5.2) can be rewritten as

$$\begin{aligned} H_1(T) &= TB'A' - (q^2B' + \alpha I)TA' - (qA' + v(q-1)I)TB' \\ &\quad + (q^3A'B' + \alpha qA' + vq^2(q-1)B' + v\alpha(q-1)I)T. \end{aligned} \quad (5.3)$$

Next, let $T \in \mathcal{B}_1$ such that $TA'x = A'x$ and $Tx = qx$; then $H_1(T)x = \alpha(v - \beta)q(q-1)x$ and so $\beta = v$ since $\alpha = \mu(q+1)(q-1) \neq 0$. Thus we have $B'A'x = \alpha(A'x + vx)$. Now let $T \in \mathcal{B}_1$ such that $TA'y = -qA'y$, $TB'y = \alpha y$ and $Ty = 0$. Then $H_1(T)y = -\alpha v(q-1)y$ and so $\alpha v(q-1) = 0$. Thus we have (1) $v = 0$. Hence $B'A'x = \alpha A'x$ and (5.3) is reduced to

$$H_1(T) = TB'A' - (q^2B' + \alpha I)TA' - qA'TB' + (q^3A'B' + \alpha qA')T. \quad (5.4)$$

Let $z \in X$ with $B'z = 0$. If $B'A'z \neq 0$, we have also $B'A'z = \alpha A'z$ as shown above with z in place of x . Suppose $B'A'z = 0$; then $B'(x+z) = 0$ but $B'A'(x+z) = B'A'x \neq 0$, and so $B'A'(x+z) = \alpha A'(x+z)$ whence $B'A'z = \alpha A'z$. In either case, we have (2) $B'A'z = \mu(q+1)(q-1)A'z$ for all $z \in X$ with $B'z = 0$.

It remains to show that $A'^2x = 0$. Note that $A'x \neq 0$ since $B'A'x \neq 0$. Suppose first that A'^2x and $A'x$ are \mathbb{C} -dependent and write $A'^2x = \theta A'x$ for some $\theta \in \mathbb{C}$. Let $T \in \mathcal{B}_1$ such that $TA'x = A'x$. Then $TA'^2x = \theta A'x$, $TB'A'x = \alpha A'x$ and $TB'A'^2x = \alpha \theta A'x$. From (5.4) it follows that $H_1(T)A'x = \alpha \theta q^2(q-1)A'x$. So $\theta = 0$ and then $A'^2x = 0$, as desired. Suppose next that A'^2x and $A'x$ are \mathbb{C} -independent.

Then $B'A'^2x \in \text{span}\{A'^2x, A'x\}$; for otherwise there would exist $T \in B_1$ such that $TB'A'^2x = A'x$ and $TA'^2x = 0$, and so $H_1(T)A'x = A'x$ by (5.4), a contradiction. Hence we may write $B'A'^2x = \ell A'^2x + kA'x$ for some $\ell, k \in \mathbb{C}$. Let $T \in B_1$ such that $TA'^2x = A'x$ and $TA'x = 0$. Then $TB'A'x = 0$ and $H_1(T)A'x = (\ell - \alpha(1 + q^2))A'x$ by (5.4). Thus $\ell = \alpha(1 + q^2)$.

If $A'^2x, A'x, x$ were \mathbb{C} -independent, then there would exist $T \in B_1$ such that $TA'^2x = x, TA'x = 0$ and $Tx = qx$; then $TB'A'(A'x + x) = \ell x = \alpha(1 + q^2)x, TA'(A'x + x) = x, TB'(A'x + x) = 0$ and $T(A'x + x) = qx$, and so $H_1(T)(A'x + x) = \alpha q^2(A'x + x)$ by (5.4), a contradiction. Hence, we may write $A'^2x = mA'x + nx$, where $m, n \in \mathbb{C}$. Then $B'A'^2x = \alpha mA'x = \ell A'^2x + kA'x$ and so the \mathbb{C} -independence of A'^2x and $A'x$ implies that $\ell = \alpha(1 + q^2) = 0$ and $k = \alpha m$. Thus $q^2 = -1$. Let $T \in B_1$ such that $TA'^2x = -qnx$ and $TA'x = A'x$. Then $TB'A'^2x = \alpha mA'x$ and $TB'A'x = \alpha A'x$, and so $H_1(T)A'x = \alpha m(1 - q)A'x$ by (5.4). Thus $m = 0$ and so $B'A'^2x = 0$ and $A'^2x = nx$. Let $\gamma \in \mathbb{C}$ with $\gamma^2 = -n/q$. Let $T \in B_1$ such that $TA'x = 0$ and $Tx = x$. Then $TB'A'(A'x + \gamma x) = TB'(A'x + \gamma x) = 0, TA'(A'x + \gamma x) = nx$ and $T(A'x + \gamma x) = \gamma x$. Thus $H_1(T)(A'x + \gamma x) = \alpha \gamma q(A'x + \gamma x)$ by (5.4), implying that $\gamma = 0$. Hence $n = 0$ and so (3) $A'^2x = 0$ follows as desired. \square

Lemma 5.6. If $B'A'^2y \neq 0$, then (1) $v = 0$, (2) $B'A'^2y = \mu(q + 1)(q - 1)A'^2y$, (3) $A'^3y = 0$, (4) $A'^2y \in \text{span}\{A'y, B'y, y\}$, (5) $B'^2y \in \text{span}\{A'y, B'y, y\}$, (6) $A'B'y \neq 0$.

Proof. Since $B'(A'y) = 0$ but $B'A'(A'y) \neq 0$, (1), (2) and (3) follow from Lemma 5.5.

Now we show that $A'^2y \in \text{span}\{A'y, B'y, y\}$. Assume on the contrary that $A'^2y, A'y, B'y, y$ are \mathbb{C} -independent. Let $T \in B_1$ such that $TA'^2y = Ty = A'y + y, TA'y = -(A'y + y)$ and $TB'y = 0$. Then $TB'A'(A'y + y) = \alpha(A'y + y)$, where $\alpha = \mu(q + 1)(q - 1) \neq 0$, and $TA'(A'y + y) = TB'(A'y + y) = T(A'y + y) = 0$, and so $H_1(T)(A'y + y) = \alpha(A'y + y)$ by (5.4), a contradiction. Therefore (4) $A'^2y \in \text{span}\{A'y, B'y, y\}$ follows.

Thus we may write $A'^2y = \xi A'y + \eta B'y + \zeta y$ for some $\xi, \eta, \zeta \in \mathbb{C}$. Since $A'^3y = 0$ but $A'^2y \neq 0$, then $A'^2y, A'y$ and y are \mathbb{C} -independent, and so $\eta \neq 0$. Comparing $B'A'^2y = \alpha A'^2y$ with $B'A'^2y = B'(\xi A'y + \eta B'y + \zeta y) = \eta B'^2y + \zeta B'y$, we obtain that $B'^2y \in \text{span}\{A'^2y, B'y\} \subseteq \text{span}\{A'y, B'y, y\}$ and so (5) holds.

From $0 = A'^3y = A'(\xi A'y + \eta B'y + \zeta y)$, it follows that $A'B'y = -\eta^{-1}(\xi A'^2y + \zeta A'y)$. Assume on the contrary $A'B'y = 0$. Then $\xi = \zeta = 0$, that is, $A'^2y = \eta B'y$, and so $B'A'^2y = \eta B'^2y$. On the other hand, $B'A'^2y = \alpha A'^2y = \alpha \eta B'y$, so $B'^2y = \alpha B'y$. Let $T \in B_1$ such that $TA'y = y$ and $TB'y = Ty = 0$. Then $TA'(y + \alpha^{-1}q^2B'y) = y$ and $TB'A'(y + \alpha^{-1}q^2B'y) = TB'(y + \alpha^{-1}q^2B'y) = T(y + \alpha^{-1}q^2B'y) = 0$. Thus $H_1(T)(y + \alpha^{-1}q^2B'y) = -\alpha(y + \alpha^{-1}q^2B'y)$ by (5.4), a contradiction. Therefore (6) $A'B'y \neq 0$ follows. \square

Since $B'A'x = BAx - \mu Ax - \nu Bx + \mu \nu x$, it follows from Lemma 5.3 that $B'A'x \in \text{span}\{Ax, Bx, x\} = \text{span}\{A'x, B'x, x\}$ for all $x \in X$. Moreover, we have

Lemma 5.7. $B'A'x \in \text{span}\{A'y, B'y, y\}$ for all $x \in X$.

Proof. Assume on the contrary that $B'A'x, A'y, B'y, y$ are \mathbb{C} -independent for some $x \in X$, then by [5, Lemma 2.1] there exists a nonzero $\lambda \in \mathbb{C}$ such that $B'A'x, A'y + \lambda A'x, B'y + \lambda B'x$ and $y + \lambda x$ are \mathbb{C} -independent. As $B'A'(y + \lambda x) = \lambda B'A'x$, it follows that $B'A'(y + \lambda x), A'(y + \lambda x), B'(y + \lambda x)$ and $y + \lambda x$ are \mathbb{C} -independent, contradicting Lemma 5.3. Hence, $B'A'x \in \text{span}\{A'y, B'y, y\}$ for every $x \in X$. \square

Lemma 5.8. Either $B'A' = 0$ or $\dim X \leq 3$.

Proof. As a vector space cannot be the set-theoretic union of two proper subspaces, it suffices to show that, for each $x \in X$, either $B'A'x = 0$ or $x \in \text{span}\{A'y, B'y, y\}$. Let $x \in X$ such that $B'A'x \neq 0$. By

Lemma 5.3, for every $\xi \in \mathbb{C}$, there exist $\mu_\xi, v_\xi \in \mathbb{C}$ such that $(B - \mu_\xi I)(A - v_\xi I)(x + \xi y) = 0$. From $(B - \mu_\xi I)(A - v_\xi I) = (B' - (\mu_\xi - \mu)I)(A' - (v_\xi - v)I)$, it follows that

$$\begin{aligned} B'A'x &= (\mu_\xi - \mu)A'x + (v_\xi - v)B'x - (\mu_\xi - \mu)(v_\xi - v)x \\ &\quad + \xi((\mu_\xi - \mu)A'y + (v_\xi - v)B'y - (\mu_\xi - \mu)(v_\xi - v)y). \end{aligned} \quad (5.5)$$

We prove first some properties about the coefficients μ_ξ, v_ξ corresponding to ξ .

(1) If $\mu_{\xi_1} = \mu_{\xi_2}$ and $v_{\xi_1} = v_{\xi_2}$ for $\xi_1, \xi_2 \in \mathbb{C}$, then $\xi_1 = \xi_2$.

Assume on the contrary $\xi_1 \neq \xi_2$. The difference of the expressions with ξ_1 and ξ_2 , respectively, in place of ξ in (5.5) yields

$$(\xi_1 - \xi_2)((\mu_{\xi_1} - \mu)A'y + (v_{\xi_1} - v)B'y - (\mu_{\xi_1} - \mu)(v_{\xi_1} - v)y) = 0.$$

Then the \mathbb{C} -independence of $A'y, B'y$ and y implies $\mu_{\xi_1} - \mu = v_{\xi_1} - v = 0$, and so $B'A'x = 0$ by (5.5), a contradiction. Hence, $\xi_1 = \xi_2$.

(2) If $v_{\xi_1} = v_{\xi_2} = v$ or $\mu_{\xi_1} = \mu_{\xi_2} = \mu$ for $\xi_1, \xi_2 \in \mathbb{C}$, then $\xi_1 = \xi_2$.

Assume on the contrary $\xi_1 \neq \xi_2$. Suppose first $v_{\xi_1} = v_{\xi_2} = v$; then $\mu_{\xi_1} \neq \mu_{\xi_2}$ by (1). It follows from (5.5) that

$$B'A'x = (\mu_{\xi_1} - \mu)(A'x + \xi_1 A'y) = (\mu_{\xi_2} - \mu)(A'x + \xi_2 A'y),$$

which implies $A'x \in \mathbb{C}A'y$. From $B'A'y = 0$, it follows that $B'A'x = 0$, a contradiction.

Suppose next $\mu_{\xi_1} = \mu_{\xi_2} = \mu$; then $v_{\xi_1} \neq v_{\xi_2}$ by (1). It follows from (5.5) that

$$B'A'x = (v_{\xi_1} - v)(B'x + \xi_1 B'y) = (v_{\xi_2} - v)(B'x + \xi_2 B'y),$$

which implies that $B'x = \omega B'y$ for some $\omega \in \mathbb{C}$ and so $B'A'x = \tau B'y$ for some nonzero $\tau \in \mathbb{C}$. Now $B'(x - \omega y) = 0$ but $B'A'(x - \omega y) = B'A'x \neq 0$, so we have by Lemma 5.5 that (i) $v = 0$, (ii) $B'A'^2y = \alpha A'^2y$, (iii) $B'A'(x - \omega y) = \alpha A'(x - \omega y)$ and (iv) $A'^2(x - \omega y) = 0$, where $\alpha = \mu(q+1)(q-1) \neq 0$. From (iii) and $B'A'x = \tau B'y$, it follows that $\tau B'y = \alpha A'(x - \omega y)$ and so $A'B'y = 0$ by (iv). Then $B'A'^2y = 0$ by Lemma 5.6, and so $A'^2y = 0$ by (ii). Let $T \in \mathcal{B}_1$ such that $TA'y = y$; then $H_1(T)A'y = \alpha q A'y$ by (5.3), a contradiction. Therefore, $\xi_1 = \xi_2$ in either situation.

(3) If $\mu_{\xi_0} = \mu_{\xi_1} = \mu_{\xi_2}$ or $v_{\xi_0} = v_{\xi_1} = v_{\xi_2}$ for $\xi_0, \xi_1, \xi_2 \in \mathbb{C}$, then ξ_0, ξ_1, ξ_2 are not all distinct.

Consider first the situation that $\mu_{\xi_0} = \mu_{\xi_1} = \mu_{\xi_2}$. The difference of the expressions with ξ_i and ξ_j , respectively, in place of ξ in (5.5) yields

$$\begin{aligned} (v_{\xi_i} - v_{\xi_j})(B'x - (\mu_{\xi_0} - \mu)x) &= (\xi_j - \xi_i)(\mu_{\xi_0} - \mu)A'y \\ &\quad + (\xi_j(v_{\xi_j} - v) - \xi_i(v_{\xi_i} - v))B'y - (\mu_{\xi_0} - \mu)(\xi_j(v_{\xi_j} - v) - \xi_i(v_{\xi_i} - v))y \end{aligned}$$

for all $i, j = 0, 1, 2$. If ξ_0, ξ_1, ξ_2 are all distinct, then so are $v_{\xi_0}, v_{\xi_1}, v_{\xi_2}$ by (1). Hence,

$$\begin{aligned} B'x - (\mu_{\xi_0} - \mu)x &= (v_{\xi_i} - v_{\xi_0})^{-1}((\xi_0 - \xi_i)(\mu_{\xi_0} - \mu)A'y + (\xi_0(v_{\xi_0} - v) \\ &\quad - \xi_i(v_{\xi_i} - v))B'y - (\mu_{\xi_0} - \mu)(\xi_0(v_{\xi_0} - v) - \xi_i(v_{\xi_i} - v))y) \end{aligned} \quad (5.6)$$

for $i = 1, 2$. Comparing the expressions with $i = 1$ and $i = 2$, respectively, in (5.6), we obtain

$$(v_{\xi_1} - v_{\xi_0})^{-1}(\xi_0 - \xi_1)(\mu_{\xi_0} - \mu) = (v_{\xi_2} - v_{\xi_0})^{-1}(\xi_0 - \xi_2)(\mu_{\xi_0} - \mu), \quad (5.7)$$

and

$$(v_{\xi_1} - v_{\xi_0})^{-1}(\xi_0 v_{\xi_0} - \xi_1 v_{\xi_1} - v(\xi_0 - \xi_1)) = (v_{\xi_2} - v_{\xi_0})^{-1}(\xi_0 v_{\xi_0} - \xi_2 v_{\xi_2} - v(\xi_0 - \xi_2)). \quad (5.8)$$

By (2) we have $\mu_{\xi_0} - \mu \neq 0$ and so (5.7) is reduced to

$$(v_{\xi_1} - v_{\xi_0})^{-1}(\xi_0 - \xi_1) = (v_{\xi_2} - v_{\xi_0})^{-1}(\xi_0 - \xi_2). \quad (5.9)$$

From (5.8) and (5.9) it follows that

$$(v_{\xi_1} - v_{\xi_0})^{-1}(\xi_0 v_{\xi_0} - \xi_1 v_{\xi_1}) = (v_{\xi_2} - v_{\xi_0})^{-1}(\xi_0 v_{\xi_0} - \xi_2 v_{\xi_2}),$$

or equivalently

$$(\xi_0(v_{\xi_0} - v_{\xi_1}) + v_{\xi_1}(\xi_0 - \xi_1))(v_{\xi_2} - v_{\xi_0}) = (\xi_0(v_{\xi_0} - v_{\xi_2}) + v_{\xi_2}(\xi_0 - \xi_2))(v_{\xi_1} - v_{\xi_0}),$$

and so

$$v_{\xi_1}(\xi_0 - \xi_1)(v_{\xi_2} - v_{\xi_0}) = v_{\xi_2}(\xi_0 - \xi_2)(v_{\xi_1} - v_{\xi_0}).$$

From the last identity and (5.9), it follows that $(\xi_0 - \xi_1)(v_{\xi_1} - v_{\xi_2})(v_{\xi_0} - v_{\xi_2}) = 0$, a contradiction. Therefore, ξ_0, ξ_1, ξ_2 are not all distinct. The proof for the situation $v_{\xi_0} = v_{\xi_1} = v_{\xi_2}$ is similar.

By (1)–(3), we may choose $\xi_1, \xi_2 \in \mathbb{C}$ with $\xi_1 \neq \xi_2$ such that v_{ξ_1}, v_{ξ_2}, v are all distinct and also that $\mu_{\xi_1}, \mu_{\xi_2}, \mu$ are all distinct. Since $B'A'x \in \text{span}\{A'y, B'y, y\}$ by Lemma 5.7, it follows from (5.5) that

$$(\mu_{\xi_1} - \mu)A'x + (v_{\xi_1} - v)B'x - (\mu_{\xi_1} - \mu)(v_{\xi_1} - v)x \in \text{span}\{A'y, B'y, y\}$$

and

$$(\mu_{\xi_2} - \mu)A'x + (v_{\xi_2} - v)B'x - (\mu_{\xi_2} - \mu)(v_{\xi_2} - v)x \in \text{span}\{A'y, B'y, y\}.$$

Eliminating $A'x$ and $B'x$, respectively, from the above two relations, we get

$$\Delta_{\xi_1, \xi_2} B'x - (\mu_{\xi_1} - \mu)(\mu_{\xi_2} - \mu)(v_{\xi_2} - v_{\xi_1})x \in \text{span}\{A'y, B'y, y\} \quad (5.10)$$

and

$$\Delta_{\xi_1, \xi_2} A'x - (v_{\xi_1} - v)(v_{\xi_2} - v)(\mu_{\xi_1} - \mu_{\xi_2})x \in \text{span}\{A'y, B'y, y\}, \quad (5.11)$$

where $\Delta_{\xi_1, \xi_2} = (\mu_{\xi_1} - \mu)(v_{\xi_2} - v) - (\mu_{\xi_2} - \mu)(v_{\xi_1} - v)$. Clearly, if $\Delta_{\xi_1, \xi_2} = 0$, then $x \in \text{span}\{A'y, B'y, y\}$ as desired. If $\Delta_{\xi_1, \xi_2} \neq 0$, then $A'x - \alpha_0 x, B'x - \beta_0 x \in \text{span}\{A'y, B'y, y\}$ for some nonzero $\alpha_0, \beta_0 \in \mathbb{C}$. From $A'x - \alpha_0 x \in \text{span}\{A'y, B'y, y\}$, it follows that $B'A'x - \alpha_0 B'x \in \text{span}\{B'^2 y, B'y\}$ and so $B'x \in \text{span}\{B'^2 y, A'y, B'y, y\}$. Hence $x \in \text{span}\{B'^2 y, A'y, B'y, y\}$ follows from $B'x - \beta_0 x \in \text{span}\{A'y, B'y, y\}$. If $B'A'^2 y \neq 0$, then $B'^2 y \in \text{span}\{A'y, B'y, y\}$ by Lemma 5.6 and consequently $x \in \text{span}\{A'y, B'y, y\}$ as desired.

So we assume $B'A'^2 y = 0$. Set $x' = B'y$. Suppose first $B'A'x' \neq 0$. Proceed as before with x, μ_{ξ}, v_{ξ} replaced by x', μ'_{ξ}, v'_{ξ} . We claim that there exist ξ_1, ξ_2 such that $\mu'_{\xi_1}, \mu'_{\xi_2}, \mu$ are all distinct, $v'_{\xi_1}, v'_{\xi_2}, v$ are all distinct and $\Delta'_{\xi_1, \xi_2} = (\mu'_{\xi_1} - \mu)(v'_{\xi_2} - v) - (\mu'_{\xi_2} - \mu)(v'_{\xi_1} - v) \neq 0$. Assume on the contrary $\Delta'_{\xi_i, \xi_j} = 0$ for all such pair ξ_i, ξ_j . From the $'$ -version of (5.5) we obtain

$$\begin{aligned} (v'_{\xi_j} - v)B'A'x' &= (v'_{\xi_j} - v)((\mu'_{\xi_i} - \mu)A'x' + (v'_{\xi_i} - v)B'x' - (\mu'_{\xi_i} - \mu)(v'_{\xi_i} - v)x' \\ &\quad + \xi_i((\mu'_{\xi_i} - \mu)A'y + (v'_{\xi_i} - v)B'y - (\mu'_{\xi_i} - \mu)(v'_{\xi_i} - v)y)) \end{aligned}$$

and

$$\begin{aligned} (v'_{\xi_i} - v)B'A'x' &= (v'_{\xi_i} - v)\left((\mu'_{\xi_j} - \mu)A'x' + (v'_{\xi_j} - v)B'x' - (\mu'_{\xi_j} - \mu)(v'_{\xi_j} - v)x'\right. \\ &\quad \left.+ \xi_j((\mu'_{\xi_j} - \mu)A'y + (v'_{\xi_j} - v)B'y - (\mu'_{\xi_j} - \mu)(v'_{\xi_j} - v)y)\right). \end{aligned}$$

The difference of them gives

$$\begin{aligned} (v'_{\xi_j} - v'_{\xi_i})B'A'(B'y) &= (\xi_i(\mu'_{\xi_i} - \mu)(v'_{\xi_j} - v) - \xi_j(\mu'_{\xi_j} - \mu)(v'_{\xi_i} - v))A'y \\ &\quad + (v'_{\xi_i} - v)(v'_{\xi_j} - v)((\xi_i - \xi_j) - (\mu'_{\xi_i} - \mu'_{\xi_j}))B'y \\ &\quad + (v'_{\xi_i} - v)(v'_{\xi_j} - v)(\mu(\xi_i - \xi_j) + (\xi_j\mu'_{\xi_j} - \xi_i\mu'_{\xi_i}))y. \end{aligned} \quad (5.12)$$

Choose $\xi_1, \xi_2, \xi_3 \in \mathbb{C}$ such that $v'_{\xi_1}, v'_{\xi_2}, v'_{\xi_3}, v$ are all distinct and $\mu'_{\xi_1}, \mu'_{\xi_2}, \mu'_{\xi_3}, \mu$ are all distinct too. Set $i = 1$ and $j = 2$, respectively, in (5.12). Comparing the corresponding coefficients of the resulted identities, we have

$$(v'_{\xi_2} - v'_{\xi_1})^{-1}(\xi_1(\mu'_{\xi_1} - \mu)(v'_{\xi_2} - v) - \xi_2(\mu'_{\xi_2} - \mu)(v'_{\xi_1} - v)) \quad (5.13)$$

$$\begin{aligned} &= (v'_{\xi_3} - v'_{\xi_1})^{-1}(\xi_1(\mu'_{\xi_1} - \mu)(v'_{\xi_3} - v) - \xi_3(\mu'_{\xi_3} - \mu)(v'_{\xi_1} - v)), \\ (v'_{\xi_2} - v'_{\xi_1})^{-1}(v'_{\xi_2} - v)((\xi_1 - \xi_2) - (\mu'_{\xi_1} - \mu'_{\xi_2})) &\quad (5.14) \\ &= (v'_{\xi_3} - v'_{\xi_1})^{-1}(v'_{\xi_3} - v)((\xi_1 - \xi_3) - (\mu'_{\xi_1} - \mu'_{\xi_3})), \end{aligned}$$

and

$$\begin{aligned} (v'_{\xi_2} - v'_{\xi_1})^{-1}(v'_{\xi_2} - v)(\mu(\xi_1 - \xi_2) + (\xi_2\mu'_{\xi_2} - \xi_1\mu'_{\xi_1})) &\quad (5.15) \\ &= (v'_{\xi_3} - v'_{\xi_1})^{-1}(v'_{\xi_3} - v)(\mu(\xi_1 - \xi_3) + (\xi_3\mu'_{\xi_3} - \xi_1\mu'_{\xi_1})). \end{aligned}$$

Since $(\mu'_{\xi_1} - \mu)(v'_{\xi_j} - v) = (\mu'_{\xi_j} - \mu)(v'_{\xi_1} - v)$, (5.13) is reduced to

$$(v'_{\xi_2} - v'_{\xi_1})^{-1}(v'_{\xi_2} - v)(\xi_1 - \xi_2) = (v'_{\xi_3} - v'_{\xi_1})^{-1}(v'_{\xi_3} - v)(\xi_1 - \xi_3). \quad (5.16)$$

By (5.16) it follows from (5.14) that

$$(v'_{\xi_2} - v'_{\xi_1})^{-1}(v'_{\xi_2} - v)(\mu'_{\xi_1} - \mu'_{\xi_2}) = (v'_{\xi_3} - v'_{\xi_1})^{-1}(v'_{\xi_3} - v)(\mu'_{\xi_1} - \mu'_{\xi_3}). \quad (5.17)$$

Comparing (5.16) and (5.17), we get

$$(\xi_1 - \xi_3)(\mu'_{\xi_1} - \mu'_{\xi_2}) = (\xi_1 - \xi_2)(\mu'_{\xi_1} - \mu'_{\xi_3}),$$

or equivalently

$$\xi_3(\mu'_{\xi_1} - \mu'_{\xi_2}) + \xi_1(\mu'_{\xi_2} - \mu'_{\xi_3}) + \xi_2(\mu'_{\xi_3} - \mu'_{\xi_1}) = 0. \quad (5.18)$$

Also, by (5.16) it follows from (5.15) that

$$(v'_{\xi_2} - v'_{\xi_1})^{-1}(v'_{\xi_2} - v)(\xi_2\mu'_{\xi_2} - \xi_1\mu'_{\xi_1}) = (v'_{\xi_3} - v'_{\xi_1})^{-1}(v'_{\xi_3} - v)(\xi_3\mu'_{\xi_3} - \xi_1\mu'_{\xi_1}).$$

Comparing this with (5.16), we get

$$(\xi_1 - \xi_3)(\xi_2 \mu'_{\xi_2} - \xi_1 \mu'_{\xi_1}) = (\xi_1 - \xi_2)(\xi_3 \mu'_{\xi_3} - \xi_1 \mu'_{\xi_1}),$$

or equivalently

$$\xi_1 \xi_2 (\mu'_{\xi_1} - \mu'_{\xi_2}) + \xi_2 \xi_3 (\mu'_{\xi_2} - \mu'_{\xi_3}) + \xi_3 \xi_1 (\mu'_{\xi_3} - \mu'_{\xi_1}) = 0. \quad (5.19)$$

From the trivial identity

$$(\mu'_{\xi_1} - \mu'_{\xi_2}) + (\mu'_{\xi_2} - \mu'_{\xi_3}) + (\mu'_{\xi_3} - \mu'_{\xi_1}) = 0$$

together with (5.18) and (5.19) it follows that $(\xi_1 - \xi_2)(\xi_2 - \xi_3)(\xi_3 - \xi_1) = 0$, contradicting our choice of ξ_1, ξ_2, ξ_3 . Therefore, there exist ξ_1, ξ_2 such that $\mu'_{\xi_1}, \mu'_{\xi_2}, \mu$ are all distinct, $v'_{\xi_1}, v'_{\xi_2}, v$ are all distinct and $\Delta'_{\xi_1, \xi_2} \neq 0$. Then $B'^2 y = B'x' \in \text{span}\{A'y, B'y, y\}$ by the $'$ -version of (5.10), and so $x \in \text{span}\{B'^2 y, A'y, B'y, y\} \subseteq \text{span}\{A'y, B'y, y\}$ as desired.

Suppose next that $B'A'x' = B'A'B'y = 0$. From $x \in \text{span}\{B'^2 y, A'y, B'y, y\}$, it follows that $B'A'x \in \mathbb{C}B'A'B'^2 y$ since $B'A'^2 y = 0$. Thus $B'A'B'^2 y \neq 0$. Set $x'' = B'^2 y$; then $B'A'x'' \neq 0$. Proceed as before with x, μ_ξ, v_ξ replaced by x'', μ''_ξ, v''_ξ . If there exist ξ_1, ξ_2 such that $\mu''_{\xi_1}, \mu''_{\xi_2}, \mu$ are all distinct, $v''_{\xi_1}, v''_{\xi_2}, v$ are all distinct and $\Delta''_{\xi_1, \xi_2} = (\mu''_{\xi_1} - \mu)(v''_{\xi_2} - v) - (\mu''_{\xi_2} - \mu)(v''_{\xi_1} - v) = 0$, then $B'^2 y = x'' \in \text{span}\{A'y, B'y, y\}$ by the $''$ -version of (5.10), and so $x \in \text{span}\{A'y, B'y, y\}$ as desired.

The proof will be complete if we show that the assumption $\Delta''_{\xi_1, \xi_2} \neq 0$ for all such pair ξ_1, ξ_2 will lead to a contradiction. In this case, $A'B'^2 y - \alpha_0 B'^2 y \in \text{span}\{A'y, B'y, y\}$ for some nonzero $\alpha_0 \in \mathbb{C}$ by the $''$ -version of (5.11), and so $B'A'^2 B'^2 y = \alpha_0 B'A'B'^2 y \neq 0$. Let $x''' = A'B'^2 y$; then $B'A'x''' \neq 0$. Proceed as before with x, μ_ξ, v_ξ replaced by $x''', \mu'''_\xi, v'''_\xi$. If there exist ξ_1, ξ_2 such that $\mu'''_{\xi_1}, \mu'''_{\xi_2}, \mu$ are all distinct, $v'''_{\xi_1}, v'''_{\xi_2}, v$ are all distinct and $\Delta'''_{\xi_1, \xi_2} = (\mu'''_{\xi_1} - \mu)(v'''_{\xi_2} - v) - (\mu'''_{\xi_2} - \mu)(v'''_{\xi_1} - v) = 0$, then $x''' \in \text{span}\{A'y, B'y, y\}$ by the $'''$ -version of (5.10), and so $B'A'x''' = 0$, a contradiction. Hence $\Delta'''_{\xi_1, \xi_2} \neq 0$ for all such pair ξ_1, ξ_2 . So by the $'''$ -version of (5.10) we have $B'A'B'^2 y - \beta_0 A'B'^2 y \in \text{span}\{A'y, B'y, y\}$ for some nonzero $\beta_0 \in \mathbb{C}$. Since $B'A'B'^2 y \in \text{span}\{A'y, B'y, y\}$ by Lemma 5.7, it follows that $A'B'^2 y \in \text{span}\{A'y, B'y, y\}$, and so $B'A'^2 B'^2 y = 0$, a contradiction again. \square

6. Proof of Theorem 5.2

As we have seen before, if $\dim X \leq 3$ then $L(X) \simeq M_3(\mathbb{C})$ and so $H(T)^3 = 0$ follows from the quasi-nilpotency of $H(T)$ for all $T \in \mathcal{B}$. So we assume $\dim X \geq 4$ and hence $B'A' = 0$ by Lemma 5.8. In this case (5.2) can be rewritten as

$$\begin{aligned} H_1(T) = & -(q^2 B' + \alpha I)TA' - (qA' + \nu(q-1)I)TB' \\ & + (q^3 A'B' + \alpha qA' + \nu q^2(q-1)B' + \alpha \nu(q-1)I)T \end{aligned} \quad (6.1)$$

for $T \in \mathcal{B}_1$, where $\alpha = \mu(q+1)(q-1)$. We claim that it suffices to consider the case $\alpha = \nu(q-1) = 0$. We proceed according as whether A'^2 and A' are locally dependent.

Case 1. $A'^2 x$ and $A'x$ are \mathbb{C} -independent for some $x \in X$.

Let $T \in \mathcal{B}_1$ such that $TA'^2 x = A'x$ and $TA'x = 0$. Then $H_1(T)A'x = -\alpha A'x$ by (6.1), implying $\alpha = 0$ as desired. So (6.1) is reduced to

$$H_1(T) = -q^2 B'TA' - (qA' + \nu(q-1)I)TB' + (q^3 A'B' + \nu q^2(q-1)B')T \quad (6.2)$$

for $T \in \mathcal{B}_1$. Let $C = qA' + \nu(q-1)I$. Then $A'C = CA' = qA'^2 + \nu(q-1)A'$, $B'C = \nu(q-1)B'$ and $CB' = qA'B' + \nu(q-1)B'$. Then (6.2) can be simplified as

$$H_1(T) = -q^2B'TA' - CTB' + q^2CB'T \quad (6.3)$$

for $T \in \mathcal{B}_1$.

Now we proceed to show either $\nu(q-1) = 0$ or $H_1(T)^3 = 0$ for $T \in \mathcal{B}_1$. Assume on the contrary $\nu(q-1) \neq 0$. Consider first the situation $A'B' = 0$. Then $CB' = B'C = \nu(q-1)B'$ and so

$$H_1(T) = -CTB' + q^2B'T(\nu(q-1)I - A').$$

If $B'Cz \notin \mathbb{C}(\nu(q-1)I - A')Cz$ for some $z \in X$, there would exist $T \in \mathcal{B}_1$ such that $TB'Cz = z$ and $T(\nu(q-1)I - A')Cz = 0$, then $H_1(T)Cz = -Cz$, a contradiction. Hence $B'Cx \in \mathbb{C}(\nu(q-1)I - A')Cx$ for all $x \in X$. By [7, Lemma 2.7], $B'C = \beta(\nu(q-1)I - A')C$ for some nonzero $\beta \in \mathbb{C}$. Thus $\nu(q-1)B'^2 = B'^2C = \beta B'(\nu(q-1)I - A')C = \beta\nu(q-1)B'C$ and so $B'^2 = \beta B'C$. Let $x \in X$ such that $B'x \neq 0$ and let $T \in \mathcal{B}_1$ such that $TB'x = B'x$. Then $H_1(T)B'x = \beta(\nu(q-1))^2(q^2-1)B'x$. This means $q+1=0$ as $\nu(q-1) \neq 0$. Thus $C = qA' + \nu(q-1)I = \nu(q-1)I - A'$. From $B'C = \beta(\nu(q-1)I - A')C$ it follows that $B'C = \beta C^2$. Using $B'C = CB' = \beta C^2 = \beta^{-1}B'^2 = \nu(q-1)B'$, $q = -1$ and $H_1(T) = -CTB' + B'TC$, we readily verify $H_1(T)^3 = 0$ for all $T \in \mathcal{B}_1$ and so we are done.

Consider next the situation $A'B' \neq 0$. If $A'B'x \notin B'X$ for some $x \in X$, there would exist $T \in \mathcal{B}_1$ such that $TA'B'x = x$ and $TB'x = TB'^2x = 0$, and so $H_1(T)B'x = -q^2B'x$ by (6.3), a contradiction. Thus $A'B'X \subseteq B'X$ and then $CB'X \subseteq B'X$. In view of (6.3) we have $H_1(T)X \subseteq B'X + CTB'X$. Thus if $\text{rank}(B') \leq 1$, then $\text{rank}H_1(T) \leq 2$ for each $T \in \mathcal{B}_1$. Then $H_1(T)$ is nilpotent, whence $H_1(T)^3 = 0$ for all $T \in \mathcal{B}_1$ and so we are done. Hence we may assume $\text{rank}(B') \geq 2$.

If $A'Cz$, $B'Cz$ and Cz are \mathbb{C} -independent for some $z \in X$, then there would exist $T \in \mathcal{B}_1$ such that $TA'Cz = TCz = 0$ and $TB'Cz = z$, and so $H_1(T)Cz = -Cz$ by (6.3), a contradiction. Hence $A'Cz$, $B'Cz$ and Cz are \mathbb{C} -dependent for any $z \in X$. By [5, Theorem 2.4], there are three cases to be investigated.

Subcase 1. There exists a subspace V of X with $\dim V \leq 3$ such that $A'CX \cup B'CX \cup CX \subseteq V$.

Since $B'C = \nu(q-1)B' \neq 0$, so $B'X = B'CX \subseteq V$, $\text{rank}(B') \leq \text{rank}(C)$ and B' maps CX onto $B'CX = B'X$. The \mathbb{C} -independence of A'^2x and $A'x$ implies $A'Cx = qA'^2x + \nu(q-1)A'x \neq 0$. Since $B'A'Cx = 0$ and $A'Cx = CA'x \in CX$, the kernel of $B' : CX \rightarrow B'X$ is not 0. Hence $\text{rank}(B') < \text{rank}(C)$. Then $\text{rank}(B') = 2$ and $\text{rank}(C) = 3$, that is $CX = V$. Thus $\text{kernel}(B') \cap V$ has dimension 1. Since $B'A'C = 0$, we have $\dim(A'CX) \leq 1$. Recall that $A'B'X \subseteq B'X$. From $0 \neq A'B'X \subseteq A'V = A'CX$, it follows that $A'CX = A'B'X \subseteq B'X$. Now $\text{rank}(A'C) = 1$ and $\text{rank}(B'C) = \text{rank}(B) = 2$, so $A'C$ and $B'C$ are not locally \mathbb{C} -dependent by [5, Theorem 2.8]. That is, there exists $x_0 \in X$ such that $A'Cx_0$ and $B'Cx_0$ are \mathbb{C} -independent. Since $A'Cx_0$, $B'Cx_0$ and Cx_0 are \mathbb{C} -dependent, $Cx_0 \in \text{span}\{A'Cx_0, B'Cx_0\} \subseteq A'CX + B'CX \subseteq B'X$. For any $z \in X$, there exists a nonzero $\lambda \in \mathbb{C}$ such that $A'C(x_0 + \lambda z)$ and $B'C(x_0 + \lambda z)$ are \mathbb{C} -independent by [5, Lemma 2.1]. As above, the \mathbb{C} -dependence of $A'C(x_0 + \lambda z)$, $B'C(x_0 + \lambda z)$ and $C(x_0 + \lambda z)$ yields $C(x_0 + \lambda z) \in B'X$ and so $Cz \in B'X$ for any $z \in X$. Thus $CX \subseteq B'X$, a contradiction.

Subcase 2. There exist $\theta_1, \theta_2, \theta_3 \in \mathbb{C}$, not all zero, such that $\theta_1A'C + \theta_2B'C + \theta_3C = 0$.

That is,

$$\theta_1(qA'^2 + \nu(q-1)A') + \theta_2\nu(q-1)B' + \theta_3(qA' + \nu(q-1)I) = 0. \quad (6.4)$$

If $\theta_2 = 0$, then, applying B' from the left to (6.4), we get $\theta_3\nu(q-1)B' = 0$. Hence $\theta_3 = 0$ and so $qA'^2 + \nu(q-1)A' = 0$, contradicting the fact that A'^2x and $A'x$ are \mathbb{C} -independent. Therefore $\theta_2 \neq 0$ and then $B' \in \text{span}\{A'^2, A', I\}$ by (6.4). So $A'B' = B'A' = 0$, a contradiction.

Subcase 3. There exists an idempotent $E \in L(X)$ of rank one such that $\text{span}\{(I-E)A'C, (I-E)B'C, (I-E)C\}$ has dimension 1.

Since $C = qA' + \nu(q-1)I = qA - \nu I$, we may write $A = \gamma I + A_1$, where $\gamma = \nu q^{-1}$ and $A_1 = q^{-1}C$. If $(I-E)C = 0$, then $A_1 = q^{-1}EC$ has rank at most 1. So $H(T) = P[P[T, B], A] = P[P[T, B], A_1]$ has rank at most 2. Thus $H(T)^3 = 0$ for all $T \in \mathcal{B}$ and we are done. Assume $(I-E)C \neq 0$; then $(I-E)B'C = \theta(I-E)C$ for some $\theta \in \mathbb{C}$. If $\theta = 0$, then $B' = (\nu(q-1))^{-1}B'C = (\nu(q-1))^{-1}EB'C$ has rank at most 1, contrary to our assumption. Hence, $\theta \neq 0$. Then $(I-E)A'C = (I-E)CA' = \theta^{-1}(I-E)B'CA' = \theta^{-1}(I-E)B'A'C = 0$. Thus $A'C = EA'C$ and so $A'CX \subseteq EX$. Since $0 \neq A'Cx \in A'CX$ and EX has dimension 1, we have $A'CX = EX$. From $A'^2C = A'CA' = EA'CA'$, it follows that $A'EX = A'^2CX \subseteq EX$. Now $B'C = \theta C + E(B'C - \theta C)$, so $A'B'X = A'B'CX \subseteq A'CX + A'EX \subseteq EX$. Hence $A'B'X = EX$ and $A'B'$ has rank 1. Then $B'E = 0$ since $B'EX = B'A'B'X = 0$, and so $B'^2C = B'(\theta C + E(B'C - \theta C)) = \theta B'C$. From $B'C = \nu(q-1)B'$ it follows that $B'^2 = (\nu(q-1))^{-1}B'^2C = (\nu(q-1))^{-1}\theta B'C = \theta B'$. If $A'B'z \notin \mathbb{C}B'z$ for some $z \in X$, there would exist $T \in \mathcal{B}_1$ such that $TA'B'z = z$ and $TB'z = 0$; then $TB'^2z = 0$ and $H_1(T)B'z = -q^2B'z$ by (6.3), a contradiction. Therefore $A'B'z \in \mathbb{C}B'z$ for all $z \in X$, so $A'B' = \lambda B'$ for some nonzero $\lambda \in \mathbb{C}$ by [7, Lemma 2.7]. Thus B' has rank 1, contrary to our assumption.

Case 2. A'^2z and $A'z$ are \mathbb{C} -dependent for all $z \in X$.

In this case, $A'^2 = \gamma A'$ for some $\gamma \in \mathbb{C}$ by [7, Lemma 2.7]. Assume first $\gamma \neq 0$. Since $A'y, B'y$ and y are \mathbb{C} -independent, so are $(A' - \gamma I)y$ and $B'(A' - \gamma I)y$. Let $T \in \mathcal{B}_1$ such that $TB'(A' - \gamma I)y = (A' - \gamma I)y$ and $T(A' - \gamma I)y = 0$. Then $H_1(T)(A' - \gamma I)y = -\nu(q-1)(A' - \gamma I)y$ by (6.1), implying $\nu(q-1) = 0$ as desired. Then (6.1) is reduced to

$$H_1(T) = -(q^2B' + \alpha I)TA' - qA'TB' + qA'(q^2B' + \alpha I)T \quad (6.5)$$

for $T \in \mathcal{B}_1$. Let $T \in \mathcal{B}_1$ such that $TA'y = A'y$. Then $H_1(T)A'y = \alpha\gamma(q-1)A'y$ by (6.5) and so $\alpha = 0$, as desired.

Assume next $\gamma = 0$, that is $A'^2 = 0$. Let $T \in \mathcal{B}_1$ such that $TA'y = A'y$. Then $H_1(T)A'y = \alpha\nu(q-1)A'y$ by (6.1) and so $\alpha\nu(q-1) = 0$. Hence $\alpha = 0$ or $\nu(q-1) = 0$.

Suppose first $\alpha = 0$; then (6.1) is reduced to (6.3) where $C = qA' + \nu(q-1)I$. Let $T \in \mathcal{B}_1$ such that $TA'y = Ty = 0$ and $TB'y = y$. Then $TA'Cy = TCy = 0$ and $TB'Cy = \nu(q-1)y$, and so by (6.3) $H_1(T)Cy = -\nu(q-1)Cy$, implying $\nu(q-1) = 0$ as desired.

Suppose next $\nu(q-1) = 0$; then (6.1) is reduced to (6.5). We claim that $A'(q^2B' + \alpha I) = 0$. Otherwise, let $x \in X$ with $A'(q^2B' + \alpha I)x \neq 0$ and $T \in \mathcal{B}_1$ such that $TA'(q^2B' + \alpha I)x = x$; then $H_1(T)A'(q^2B' + \alpha I)x = qA'(q^2B' + \alpha I)x$ by (6.5), a contradiction. Hence, $A'(q^2B' + \alpha I) = 0$ and so (6.5) is reduced to

$$H_1(T) = -(q^2B' + \alpha I)TA' - qA'TB'$$

for $T \in \mathcal{B}_1$. Now it is easy to see that $H_1(T)^3 = 0$ for all $T \in \mathcal{B}_1$ since $B'A' = A'^2 = 0$ and so we are done.

Now it remains to consider the case $\alpha = \nu(q-1) = 0$. In this case, (6.1) is reduced to

$$H_1(T) = -q^2B'TA' - qA'TB' + q^3A'B'T \quad (6.6)$$

for $T \in \mathcal{B}_1$. Suppose first $A'B' = 0$; then (6.6) is reduced to

$$H_1(T) = -qA'TB' - q^2B'TA' \quad (6.7)$$

for $T \in \mathcal{B}_1$. We claim that either $A'^2 = 0$ or $B'^2 = 0$, whence $H_1(T)^3 = 0$ for all $T \in \mathcal{B}_1$. Assume on the contrary that both $A'^2 \neq 0$ and $B'^2 \neq 0$; then there exists $x \in X$ such that $A'^2x \neq 0$ and $B'^2x \neq 0$. Since $B'A' = 0$, $A'x$ and $B'x$ are \mathbb{C} -independent. If A'^2x and B'^2x are also \mathbb{C} -independent, there would exist $T \in \mathcal{B}_1$ such that $TA'^2x = x$ and $TB'^2x = qx$, and then $H_1(T)(A'x + B'x) = -q^2(A'x + B'x)$ by

(6.7), a contradiction. Hence $A'^2x = \tau B'^2x$ for some nonzero $\tau \in \mathbb{C}$. Write $\tau = \rho^2$ and $q = p^2$ for some $\rho, p \in \mathbb{C}$. Take $T \in \mathcal{B}_1$ such that $TB'^2x = x$. Then

$$H_1(T) (B'x + \rho^{-1}p^{-1}A'x) = -\rho p^3 (B'x + \rho^{-1}p^{-1}A'x),$$

by (6.7), a contradiction again. Therefore, either $A'^2 = 0$ or $B'^2 = 0$ and we are done.

Suppose next $A'B' \neq 0$. If $A'^2B' = 0$ or $A'^2B'x \notin \mathbb{C}A'B'x$ for some $x \in X$, there would exist $x \in X$ and $T \in \mathcal{B}_1$ such that $A'B'x \neq 0$, $TA'^2B'x = 0$ and $TA'B'x = x$, and then $H_1(T)A'B'x = q^3A'B'x$ by (6.6), a contradiction. Hence, $A'^2B' = \lambda A'B' \neq 0$ for some $\lambda \in \mathbb{C}$ by [7, Lemma 2.7]. If $A'B'x \notin B'X$ for some $x \in X$, there would exist $T \in \mathcal{B}_1$ such that $TA'B'x = x$ and $TB'x = TB'^2x = 0$, and then $H_1(T)B'x = -q^2B'x$ by (6.6), a contradiction. So $A'B'X \subseteq B'X$. Let $z \in X$ with $A'B'z \neq 0$. Then $A'B'z = B'z_0$ for some $z_0 \in X$ and then $A'B'z_0 = A'^2B'z = \lambda A'B'z = \lambda B'z_0$ and $B'^2z_0 = 0$. Let $T \in \mathcal{B}_1$ such that $TB'z_0 = z_0$. Then $H_1(T)B'z_0 = \lambda q^2(q-1)B'z_0$ by (6.6), implying that $q = 1$. So $H_1(T) = A'B'T - A'TB' - B'TA' = [A', [B', T]]$ for $T \in \mathcal{B}_1$. Then the conclusion of Theorem 5.2 follows from [7, Theorem 1.2].

References

- [1] Z. Abdelali, On Φ -derivations in Banach algebras, *Comm. Algebra* 34 (2006) 2437–2452.
- [2] J. Bergen, P. Grzeszczuk, Invariants of $U_q(Sl(2))$ and q -derivations, *J. Pure Appl. Algebra* 133 (1998) 27–38.
- [3] N. Boudi, On the product of derivations in Banach algebras, *Math. Proc. R. Ir. Acad.* 109 (2009) 201–212.
- [4] M. Brešar, On automorphisms of Banach algebras, *Arch. Math. (Basel)* 78 (2002) 297–302.
- [5] M. Brešar, P. Šemrl, On locally linearly dependent operators and derivations, *Trans. Amer. Math. Soc.* 351 (1999) 1257–1275.
- [6] M. Brešar, A.R. Villena, The noncommutative Singer–Werner conjecture and ϕ -derivations, *J. London Math. Soc.* 66 (2002) 710–720.
- [7] M.A. Chebotar, W.-F. Ke, P.-H. Lee, On a Brešar–Šemrl conjecture and derivations of Banach algebras, *Q. J. Math.* 57 (2006) 1–10.
- [8] C.-L. Chuang, T.-K. Lee, Algebraic q -skew derivations, *J. Algebra* 282 (2004) 1–22.
- [9] C.-L. Chuang, T.-K. Lee, q -Skew derivations and polynomial identities, *Manuscripta Math.* 116 (2005) 229–243.
- [10] C.-L. Chuang, C.-K. Liu, Extended Jacobson density theorem for rings with skew derivations, *Comm. Algebra* 35 (2007) 1391–1413.
- [11] K.R. Goodearl, Prime ideals in skew polynomial rings and quantized Weyl algebras, *J. Algebra* 150 (1992) 324–377.
- [12] K.R. Goodearl, E.R. Letzter, Prime ideals in skew and q -skew polynomial rings, *Mem. Amer. Math. Soc.* 109 (521) (1994), Amer. Math. Soc., Providence, RI.
- [13] P. Grzeszczuk, A. Leroy, J. Matczuk, Artinian properties of constants of algebraic q -skew derivations, *Israel J. Math.* 121 (2001) 265–284.
- [14] P. Grzeszczuk, J. Matczuk, The prime radical of constants of algebraic q -skew derivations, *Comm. Algebra* 30 (2002) 5331–5343.
- [15] I. Kaplansky, *Infinite Abelian Groups*, University of Michigan Press, Ann Arbor, 1954.
- [16] D.C. Kleenecke, On operator commutators, *Proc. Amer. Math. Soc.* 8 (1957) 535–536.
- [17] T.-K. Lee, C.-K. Liu, Nilpotency of q -skew derivations, *Comm. Algebra* 34 (2006) 661–669.
- [18] M. Mathieu, G.J. Murphy, Derivations mapping into radical, *Arch. Math. (Basel)* 57 (1991) 469–474.
- [19] V. Prák, Commutators in Banach algebras, *Proc. Edinburgh Math. Soc.* 22 (1979) 207–211.
- [20] C.E. Rickart, *General Theory of Banach Algebras*, Van Nostrand, New York, 1960.
- [21] F.V. Shirokov, Proof of a conjecture of Kaplansky, *Uspehi Mat. Nauk.* 11 (1956) 167–168.
- [22] A.M. Sinclair, Continuous derivations on Banach algebras, *Proc. Amer. Math. Soc.* 20 (1969) 166–170.
- [23] M.P. Thomas, Primitive ideals and derivations on noncommutative Banach algebras, *Pacific J. Math.* 159 (1993) 139–152.
- [24] Yu. V. Turovskii, V.S. Shul'man, Conditions for the massiveness of the range of a derivation of a Banach algebra and of associated differential operators, *Mat. Zametki* 42 (1987) 305–314.